



The Effect of *Jordanian Herbs* on Gut Microflora, and Immunity of Broiler Chickens

Mustafa S. AL-Rawashdeh¹

¹Department of Animal Production, Faculty of Agriculture, Mutah University, Karak 61710, Jordan.

Email: mustafa_r@mutah.edu.jo

ORCID ID: 0000-0002-7025-0692

*Corresponding author's E-mail: mustafa_r@mutah.edu.jo

Article History	Abstract
Received: 06 February 2022 Revised: 28 July 2023 Accepted: 06 August 2023	The study aimed to examine and compare the influence of adding synthetic antioxidant butylated hydroxyanisole (BHA), with two herbal medical plants (<i>Oregano germander</i>) on broiler gut microflora, and its immunity. One hundred and forty broilers were raised for three weeks and were distributed into five treatment groups. The prepared dietary treatments were included (w/w):1). Control (without supplement) 2). Germander (GER; inclusion level 1.5 %) 3). Oregano (ORE; inclusion level 2.5 %) 4). Combination of GER and ORE (CM; inclusion levels 1.5 and 2.5 %, respectively)5). BHA (0.02 %). The highest total number of ileal lactobacilli counts was found in broilers fed CM dietary treatment. On the other hand, their disgust samples (i.e. in CM) showed the lowest total count of <i>Escherichia coli</i> compared to other treatments. Among different dietary treatments, CM treatment showed the best positive effect on broiler immune system response, where it showed the highest Bursa Fabricius relative weight, and IgG values. Overall, the results showed that such dietary treatment (i.e. Combination of GER and ORE) positively enhanced gut microflora, and bird's immunity response.
CC License CC-BY-NC-SA 4.0	Keywords: Germander, Gut microflora, Ideal microflora, Oregano essential oil, Broilers.

1. Introduction

Many feed supplements are widely used in poultry diets to enhance their growth performance and enhancing their health status (Senay et al., 2019; Gopi et al., 2014; Hassan et al., 2018). Many researchers investigated the effect of adding different plant-based supplements on poultry growth performance, feed efficiency, immunity, antioxidant defense systems, and the quality of their products (Al-Rabadi et al., 2020; Al-hijazeen, 2019; Al-hijazeen et al., 2022 AL-Rawashdeh et al., 2022; Ri et al., 2017; Surai et al., 2019). Furthermore, several reasons encourage animal production scientists toward this approach; however, replacing natural with synthetic anti-oxidant/antimicrobial supplements from poultry diets was their major goal (Senay et al., 2019; Al-Hijazeen, 2019). It is well known that using synthetic additives (synthetic antioxidants and antibiotics) in animal diets became

unfavorable due to their direct and indirect side effect on human health (Gopi et al., 2014; Mohammadi Gheisar and Kim, 2018). In addition, these additives reduce poultry disease resistance (Agyare et al., 2018) for several bacterial pathogenic strains (Gopi et al., 2014; Mohammadi Gheisar and Kim, 2018). Furthermore, it had been reported that the presence of these additives in the poultry diet may be adsorbed by the animal tissues (lean meat), and so indirectly affect humans (Senay et al., 2019; Jiang and Xiong, 2016; Mohammadi Gheisar and Kim, 2018; Abbdool et al., 2017).

Nowadays, all scientists are interested in finding natural and safe alternatives, especially in the poultry feed industry (Gopi et al., 2014; Hassan et al., 2018; Mohammadi Gheisar and Kim, 2018). One of the most suggested dietary alternatives is using phytobiotics (e.g., dried medicinal plant leaves, and their extracts) (Aroche et al., 2018; Windisch et al., 2008). These plants material composed of many phenolic compounds which positively improve broiler chickens' performances, feed efficiency, nutrient digestibility, and immune function (Aroche et al., 2018; Mohammadi Gheisar and Kim, 2018; Giannenas et al., 2004; Platel and Srinivasan, (2004); Jamroz et al., 2006).

For instance, oregano, sage, green tea, and varieties of plants resources showed a positive effect on overall animal performance; enhance animal growth, health status, and final carcass quality (Botsoglou et al., 2003a, b; Attia et al., 2017; Peng et al., 2016; Ri et al., 2017; Giannenas et al. 2005). It had been reported that adding these plants supplement (as dried plants or extracts) may improve poultry feed digestibility, feed efficiency, and thus their growth performance (Jamroz et al., 2003; Botsoglou et al., 2003a; Giannenas et al. 2005; Windisch et al., 2008). The improvement in poultry performance was also connected to the positive morphological changes, increasing in digestive enzyme activity which enhances its nutrient digestibility (Li et al., 2012; Lee et al., 2003; Jang et al., 2004; Jang et al., 2007). However, it always depends on many factors such as plant components, combination used, and animal response. These beneficial effects are highly dependent on their antioxidant and antibacterial properties especially when animals have stressed conditions (Al-Hijazeen, 2019; Al-Bandak, 2007; Gopi et al., 2014; Al Bahtiti, 2012; Jaradat, 2015). In addition, it enhances the blood immune system by improving animal reducing capacity, antioxidant defense system, and the formation of free radicals at the cellular level (Park et al., 2015; Gheisar and Kim, 2018). Animal tissues may also uptake certain compounds found in herbal plants oil (extract) like the vitamin E absorption mechanism (Leal et al., 2018; Lui et al., 1995; Li et al., 2009; Botsoglou et al., 2002; 2003a; 2003b).

For instance, oregano oil (*Origanuimsyriacum* L.) extraction was analyzed by different laboratories and procedures (Ibrahim et al., 2012a; b). Its extracts are composed of various polyphenols which consider being responsible for its antioxidant activity (Giannenas et al., 2013). Volatile compounds such as *carvacrol* and *thymol* are the major antioxidant compounds in oregano oil Kosar et al., 2003; Al-Bandak, 2007; Ibrahim et al., 2012a; Kokkini, 1997). Another example is germander plant (*TeucriumPolium* L.) extract has many poly-phenolic compounds that may be useful to treat illness and prevent many human diseases (Jaradat, 2015; Mahmoudi and Nosratpour, 2013) Germander essential oil and its extracts (*wild* Jordanian: *Teucriumpolium* (Ja'adeh) were consist unique phenolic compounds that showed a strong antioxidant and radical scavenging activities (Djilas et al., 2006). *Teucrium. Polium* L. (Wild Jordanian) composition is extensively analyzed, and well documented. Jaradat (2015) summarized 50 years of scientific knowledge related to germander phytochemical composition into different categories. Furthermore, the essential oil and extracts of both plants (i.e. oregano and germander) showed significant antimicrobial effects against several pathogenic bacterial strains (Jaradat, 2015; Alhijazeen, 2018; Mohammadi Gheisar and Kim, 2018). The antimicrobial mode of action of these plants' additives was varying and inconsistent. This effect is due to differences in their functional hydroxyl or alkyl group location (Yang et al., 2015). Researchers suggested that the antimicrobial activity of these plants derived was due to their direct influence on the microflora cell membrane, ion exchange and internal protein synthesis (Burt, 2004; Ultee et al., 2002).

Both medicinal plants (Oregano and germander) are available in the wild of the south Jordanian region (Al Bahiti, 2012; Ibrahim et al., 2012a; Afifi et al., 2009) with high levels of phenolic compounds (Ibrahim et al., 2012a; Acar and Goldstein, 1996). However, many aspects could affect their potency as antioxidant/antimicrobial agents. For example, their storage condition, cultivation

period, harvesting time, and plant variety (Burt, 2004; Badi, et al., 2004; Ibrahim et al., 2012a; Afifi et al., 2009). Both *Teucrium polium* L. and *Origanum syriacum* L. are characterized by their unique volatiles and composition compared to other plant species (Ibrahim et al., 2012a; Jaradat, 2015). These alternatives are highly recommended due to many reasons, which promote the use of it as poultry feed additives (Gheisar and Kim, 2018). A little investigation had been done on their combined effect (Oregano and germander) on the broiler immune system, and their effect on gut microflora. The main purpose of this study was to examine the influence of supplementing dietary oregano and germander herbal plants alone or in combination; on the immunity and gut microflora in broiler chicken and to compare the influence of these dietary herbs with the synthetic antioxidant (BHA).

2. Materials And Methods

Birds and Diet Formulation

One hundred and forty broilers were raised for three weeks (during the growing stage; from 21 to 42 days old). The five dietary treatments of 140 broilers were randomly assigned (each treatment include 7 replicates and 4 broilers per replicate). The formulated dietary treatments were: 1). Without supplement (control); 2). Level of 1.5 % Germander (GER); 3). Level of 2.5 % Oregano (ORE); 4) Combination of GER and ORE (CM, levels of 1.5 and 2.5 %, respectively); 5). *butylated hydroxyanisole* BHA (level of 0.02 %). All dietary treatments were prepared to meet nutrient specifications suggested by poultry NRC (1994).

Immune System Response and their Organs Index

Two measurements for each replicate were reported to evaluate both spleen and *Bursa Fabricius* relative weights (in proportion to body weight %) at the age of 42 days as a good indication about the immune system response of broiler chicken by adding these plants supplements. On the final day (day 42), blood samples were taken from the jugular vein into plain vacutainer tubes, coagulated at room temperature for 10-20 mins, and then the separated serum was stored at -20°C until further analysis. Serum immunoglobulin (IgG or IgY) was measured and analyzed using a prepared kit (Chicken IgG (IgY) ELISA Life Diagnostics, Inc., Catalog Number: IGG-5) according to the manufacturer's recommendations. Samples were incubated for 5 min at 37 °C, and the absorbance (At 450 nm/after adding stop solution/within 15 min) was recorded using a microplate reader (ELISA). The concentration of IgG was calculated depending on the absorbance of the tested samples and is derived from a standard curve.

Gut Microflora

All intestine samples (two birds per replicate treatment) were cut and collected immediately after slaughtering according to Ao and Kim's (2019b) protocol with minor modifications [50]. Small intestine cut samples were vacuum packaged (Baalbaki®DZ 400/2D Vacuum Packer, 14KT6013-HZ-5), and cooled at 4°C until used. Fresh cut samples (1 g of internal digesta in duplicate) were aseptically diluted in 9 ml of 1 % sterilized peptone broth (BiolabZrt. 1141 BUDAPEST Öv u. 43, Hungary), (10-fold dilution) then the subsequent dilutions were vortex and used directly to enumerate selected digestive bacterial. This protocol evaluated the effect of adding these plant supplements on the survival of selected bacterial populations from birds DSI (Digesta of Small Intestine) at day 42. The colony-forming units (CFU/g) of certain were evaluated for all treatment groups according to the method described by Rahimi et al (2011) and Ao et al (2019). MacConkey agar (Merk, Germany) media was prepared to enumerate viable *E. coli* at 37 °C for 24 hours, and MRS (Merk, Germany) agar was used to enumerate *Lactobacillus* viable count under the anaerobic condition at 37°C for 72 hours.

Statistical Analysis: measurements were statistically examined using the procedures of a generalized linear model (Proc. GLM, SAS program, 2012). Complete randomized design (CRD) was used to analyze measurements. Tukey test or Tukey's Multiple Range test was used to determine whether there is a significant difference between the average values (the significance was determined at $P < 0.05$).

3. Results and Discussion

Gut Microflora

Ileal microflora, its varieties, stabilization, and their content of both beneficial and pathogenic were reported as effective factors affecting overall bird's health and growth performance (Song et al., 2014; Mohammadi Gheisar and Kim, 2018). Several studies (in vivo antimicrobial effect) showed that adding plants derived/phytogenic feed additives (PFAs) in broiler diet were suppress the growth of many pathogens such as *Clostridium perfringens* (Mitsch et al. 2004), *Eimeria* species (Ovied-Rondon et al., 2006; Hume et al., 2006). In Table (1), it was clearly shown that adding all feed additives suppress the growth of *E.coli* significantly compared to the control treatment. However, the CM showed the highest antibacterial effect against *E.coli* survival among the other treatments. In addition, the anti-*E.coli* growth effect of adding ORE supplements was significantly higher than adding GER and BHA alone. On the other hand, the CM showed the highest significant ($P < 0.05$) positive effect on the growth and survival of *Lactobacillus* ($6.11 \text{ Log CFU/g}^{-1}$). Oregano (ORE) additive showed a higher significant effect in increasing *Lactobacillus* count compared to GER and BHA alone. Mohammadi Gheisar and Kim (2018) reported that adding (PFAs) to the gut of poultry and swine will decrease the total count of pathogens which causes an increase in the beneficial bacteria count (*Lactobacillus*). Adding PFAs may improve the growth of some beneficial bacteria strains in the bird's digestive system, on the other hand, it may decrease the total count of certain pathogenic bacteria (Brenes et al., 2016; Mohammadi Gheisar et al., 2015a). Viveros et al. (2011) also concluded that adding GSE positively affects the total bacterial count of beneficial bacteria and suppresses the growth of some pathogenic bacteria. Researchers explain the increase in the total number of ileal *lactobacilli* counts through the ability of these bacterial strains to utilize or metabolize phenolic compounds (nutritional substances) (Garcia-Ruiz et al., 2008; Viveros et al., 2011). Tschirch (2000) reported that carvacrol (oregano and thyme main constituent) enhanced *Lactobacillus* survival and growth rate. A similar trend was found by Mohammadi Gheisar et al. (2015a) where the essential oils blend increased the *Lactobacillus* count of broilers chickens. Furthermore, phenolic compounds can increase the shedding of microbial bacteria (*Lactobacillus*) which can improve bird's immunity (Paszkiwicz et al., 2012). Other researchers documented the anti *E.coli* effect in vitro by using these plant-derived additives (Jamroz et al., 2003; 2006). Similar results found that the *E.coli* (ileal microflora) count decreased by adding herbal extract (garlic, coneflower, thyme) in broilers feed (Skyrme, 1997; Rahimi et al., 2011). Furthermore, researchers reported that the inclusion of dietary herbal plants may improve the jejunum morphology (crypt depth and villus height) and this was reflected positively in the bird's feed efficiency and gut health (Boka et al., 2014; Yang et al., 2016; Viveros et al., 2011; Samuel et al., 2017).

Table 1. Number (Log CFU/g-1Digesta) of viable Escherichia coli, Lactices inileo-cecum of broiler chickens.

Treatments	Control	Germander	Oregano	Combination	BHA	SEM
Microflora	----- Log CFU/g ¹ Digesta-----					
<i>Escherichia coli</i> ^x	7.177 ^a	4.891 ^c	4.327 ^d	4.031 ^e	6.371 ^b	0.067
<i>Lactics</i>	4.362 ^c	4.777 ^c	5.505 ^b	6.112 ^a	4.355 ^c	0.104

^x- Within a row, average values with different subscript differ significantly ($p < 0.05$).

Prepared intestinal digesta (SID) samples stored at 4 C. n=7.

Bird's Immunity Response and their Organ Index

Immunity organs relative weight (index), and IgG in blood samples of broiler fed different diets were reported (Table 2). Generally, the broiler immune system response is correlated with the immune organ size such as the spleen, thymus, and bursa (Króliczewska et al., 2017; Zhou et al., 2019). In the current study, there were no differences found among all treatments when evaluating their spleen relative weight (S. RW). However, a significant difference is identified on *Bursa Fabricious* relative weight (B. FRW) between the control and the CM treatments diets. In addition, no significant differences showed when testing B.F RW (%) among all treatment additives. The CM treatment showed higher B. F RW (%) of all birds compared to other dietary treatments which indicates a higher immunity response. However, the effect of adding various plant additives on broiler immune organs (Organ's index: such as spleen, thymus, bursa and liver) was varied among all studies done previously (Zhou et al., 2019; Ri et al., 2017; Farahat et al., 2017). This variation had been attributed to the differences between these plants derived in their effective antioxidant compounds (Abu Hafsaand

Ibrahim, 2018; Attia et al., 2017). Avian serum immunoglobulins (IgG or IgY) which are the major antibody in birds was also used to measure its immunity response (Suzuki and Lee, 2004). In this study, there was no significant difference in IgG mean level between the control diet, and the other treatment diets. In contrast, both ORE and CM treatment diets showed the highest levels (6.24 and 6.43 mg/ml, respectively). These results agreed with Hamal et al. (2006) where the normal chicken serum IgG level was in the range of 3-6 mg/ml. Several research studies reported that dietary plant-derived or their extracts in animal feed raise their serum antibody levels (Zhou et al., 2019; Farahat et al., 2017; Facina et al., 2017). Furthermore, broiler's immunity response is correlated with many factors such as type of infection, pathogens' effectiveness, and bird's genetic susceptibility factors (Oviedo-Rondon et al., 2006; Yang et al., 2016; Zahoor et al., 2018).

Table 2. The immune system response and relative weight of broilers fed different dietary supplements.

Parameter	Control	GER	ORE	CM	BHA	SEM
S. RW (%) ^x	0.114 ^a	0.116 ^a	0.127 ^a	0.128 ^a	0.122 ^a	0.005
B. F RW (%)	0.195 ^b	0.212 ^{ab}	0.217 ^{ab}	0.227 ^a	0.209 ^{ab}	0.006
IgG	5.43 ^a	5.89 ^a	6.24 ^a	6.43 ^a	5.91 ^a	0.396

Within a row, average values with different subscripts differ significantly ($p < 0.05$).

S. RW: Spleen relative weight (% of body weight).

B.F RW: Bursa Fabricius relative weight (% of body weight).

IgG: mg/ml.

N=7.

4. Conclusion

Dietary supplements affected broiler gut microflora and their immunity response. Combining both herbs treatments (i.e. CM) exposed the highest effect in enhancing bird's immunity response. In addition, it (CM) exhibited the highest effect suppressing the *E. coli* growth and enhances *Lactobacillus* survival. Their positive effect on the birds' gut microflora considers a mirror in the improvements of the bird's overall health. It is concluded that adding a combination of ORE and GER could improve the overall broiler immune response and gut microflora.

Acknowledgements:

I am deeply grateful to those who contributed to the completion of this research. I would like to thank the deanship of scientific research at Mutah University for their continuous assistance with providing access to the most recent researches in the field of study, Dr. Ghaid Alrabadi for helping in data collection and Dr. Abeer Alrawashdeh in proofreading the manuscript. Also, I would like to thank some of my colleagues in the department of Animal production at Mutah University for their invaluable support throughout the research process. Their insights and expertise were effective in the success of this project.

Conflict of interest:

The author declares no conflict of interest.

References:

- Abbood, A. A., Kassim, A. B., Jawad, H. S., Manap, Y. A., & Sazili, A. Q. (2017). Effects of feeding the herb *Borreria latifolia* on the meat quality of village chickens in Malaysia. *Poultry science*, 96(6), 1767-1782. <https://doi.org/10.3382/ps/pew460>
- Abu Hafsa, S. H., & Ibrahim, S. A. (2018). Effect of dietary polyphenol-rich grape seed on growth performance, antioxidant capacity and ileal microflora in broiler chicks. *Journal of animal physiology and animal nutrition*, 102(1), 268-275. <https://doi.org/10.1111/jpn.12688>
- Acar, J.F., and F.W. Goldstein, (1996). Disc Susceptibility test. In: Lo-Rian, V. (Ed.), *Antibiotics in laboratory medicine*. 4th Edn., William & Wilkins, Baltimore, London pp: 1-52.

- Afifi, F., Abu-Irmaileh, B., & Al-Noubani, R. (2009). Comparative analysis of the essential oils of *Teucrium polium* L. grown in different arid & semi arid habitats in Jordan. *Jordan Journal of Pharmaceutical Sciences*, 2(1).
- Agyare, C., Boamah, V. E., Zumbi, C. N., & Osei, F. B. (2018). Antibiotic use in poultry production and its effects on bacterial resistance. *Antimicrobial resistance—A global threat*, 33-51. <http://dx.doi.org/10.5772/intechopen.73725>
- ALAL-Rawashdeh, M. S., AL-Hijazeen, M. A., & AL-Rabadi, G. J. (2022). PRESERVATIVE EFFECT OF DIETARY OREGANO AND GERMANDER ON QUALITY AND STORAGE STABILITY OF RAW BROILER MEAT. *JAPS: Journal of Animal & Plant Sciences*, 32(5).
- Al-Bandak, G., & Oreopoulou, V. (2007). Antioxidant properties and composition of *Majorana syriaca* extracts. *European Journal of Lipid science and technology*, 109(3), 247-255. <https://doi.org/10.1002/ejlt.200600234>
- Al-Hijazeen, M. (2018). EVALUATE ANTI-BACTERIAL ACTIVITY OF *ROSMARINUS OFFICINALIS* LINN. EXTRACT AND *ORIGANUM SYRIACUM* L. ESSENTIAL OIL USING RAW CHICKEN MEAT. *Carpathian Journal of Food Science & Technology*, 10(2).
- Al-Hijazeen, M. (2019). Effect of *Origanum syriacum* L. essential oil on the storage stability of cooked chicken meat. *Brazilian Journal of Poultry Science*, 21(1):1-10. <http://dx.doi.org/10.1590/1806-9061-2017-0719>.
- Ao, X., and I. H. Kim. (2019a). Effects of astaxanthin produced by *Phaffiarhodozyma* on growth performance, antioxidant activities and meat quality in Pekin ducks. *Poult. Sci.* <https://doi.org/10.3382/ps/pez256>.
- Ao, X., and Kim, I. H. (2019b). Effects of grape seed extract on performance, immunity, antioxidant capacity, and meat quality in Pekin ducks. *Poultry Science*, <https://doi.org/10.1016/j.psj.2019.12.014>(Article in press).
- Aroche, R., Martinez, Y., Ruan, Z., Guan, G., Waititu, S., Nyachoti, C. M., Mas, D., and Lan, S. (2018). Dietary inclusion of a mixed powder of medicinal plant leaves enhances the feed efficiency and immune function in broiler chickens. *Journal of Chemistry*, Article ID 4073068, 6 pages, <https://doi.org/10.1155/2018/4073068>.
- Badi, H. N., Yazdani, D., Ali, S. M., & Nazari, F. (2004). Effects of spacing and harvesting time on herbage yield and quality/quantity of oil in thyme, *Thymus vulgaris* L. *Industrial crops and products*, 19(3), 231-236. <https://doi.org/10.1016/j.indcrop.2003.10.005>
- Botsoglou, N. A., Florou-Paneri, P., Christaki, E., Fletouris, D. J., & Spais, A. B. (2002). Effect of dietary oregano essential oil on performance of chickens and on iron-induced lipid oxidation of breast, thigh and abdominal fat tissues. *British poultry science*, 43(2), 223-230. <https://doi.org/10.1080/00071660120121436>
- Botsoglou, N. A., Govaris, A., Botsoglou, E. N., Grigoropoulou, S. H., & Papageorgiou, G. (2003). Antioxidant activity of dietary oregano essential oil and α -tocopheryl acetate supplementation in long-term frozen stored turkey meat. *Journal of agricultural and food chemistry*, 51(10), 2930-2936. <https://doi.org/10.1021/jf021034o>
- Botsoglou, N. A., Grigoropoulou, S. H., Botsoglou, E., Govaris, A., & Papageorgiou, G. (2003). The effects of dietary oregano essential oil and α -tocopheryl acetate on lipid oxidation in raw and cooked turkey during refrigerated storage. *Meat science*, 65(3), 1193-1200. [https://doi.org/10.1016/S0309-1740\(03\)00029-9](https://doi.org/10.1016/S0309-1740(03)00029-9)
- Brenes, A., A. Viveros, S. Chamorro, and I. Arija. (2016). Use of polyphenol-rich grape by-products in monogastric nutrition. a review. *Anim. Feed Sci. Technol.* 211:1-17.
- Burt, S. (2004). Essential oils: their antibacterial properties and potential applications in foods—a review. *International journal of food microbiology*, 94(3), 223-253. <https://doi.org/10.1016/j.ijfoodmicro.2004.03.022>
- Djilas, S. M., Markov, S. L., Cvetković, D. D., Čanadanović-Brunet, J. M., Četković, G. S., & Tumbas, V. T. (2006). Antimicrobial and free radical scavenging activities of *Teucrium montanum*. *Fitoterapia*, 77(5), 401-403. <http://dx.doi.org/10.1016/j.fitote.2006.05.019>
- Farahat, M. H., Abdallah, F. M., Ali, H. A., & Hernandez-Santana, A. (2017). Effect of dietary supplementation of grape seed extract on the growth performance, lipid profile, antioxidant status and immune response of broiler chickens. *Animal*, 11(5), 771-777. <https://doi.org/10.1017/S1751731116002251>
- Fascina, V. B., Pasquali, G. A. M., Carvalho, F. B., Muro, E. M., Vercese, F., Aoyagi, M. M., ... & Sartori, J. R. (2017). Effects of phytogenic additives and organic acids, alone or in combination, on the performance, intestinal quality and immune responses of broiler chickens. *Brazilian Journal of Poultry Science*, 19, 497-508. <https://doi.org/10.1590/1806-9061-2016-0422>
- García-Ruiz, A., Bartolomé, B., Martínez-Rodríguez, A. J., Pueyo, E., Martín-Álvarez, P. J., & Moreno-Arribas, M. V. (2008). Potential of phenolic compounds for controlling lactic acid bacteria growth in wine. *Food Control*, 19(9), 835-841. <https://doi.org/10.1016/j.foodcont.2007.08.018>

- Giannenas I, Bonos, E., Christaki, E., Florou-paneri, P. (2013). Essential oils and their applications in animal nutrition. *Med Aromatic Plants*. 2:1-12.
- Giannenas, I. A., Florou-Paneri, P., Botsoglou, N. A., Christaki, E., & Spais, A. B. (2005). Effect of supplementing feed with oregano and/or alpha-tocopheryl acetate on growth of broiler chickens and oxidative stability of meat. *Journal of Animal and Feed Sciences*, 14(3), 521. <https://doi.org/10.22358/jafs/67120/2005>
- Giannenas, I. A., Florou-Paneri, P., Papazahariadou, M., Botsoglou, N. A., Christaki, E., & Spais, A. B. (2004). Effect of diet supplementation with ground oregano on performance of broiler chickens challenged with *Eimeria tenella*. *Archiv fur Geflugelkunde*, 68(6), 247-252.
- Gopi, M., Karthik, K., Manjunathachar, H. V., Tamilmahan, P., Kesavan, M., Dashprakash, M., ... & Purushothaman, M. R. (2014). Essential oils as a feed additive in poultry nutrition. *Adv. Anim. Vet. Sci*, 2(1), 1-7. <http://www.nexusacademicpublishers.com/journal/4>.
- Hamal, K. R., Burgess, S. C., Pevzner, I. Y., & Erf, G. F. (2006). Maternal antibody transfer from dams to their egg yolks, egg whites, and chicks in meat lines of chickens. *Poultry science*, 85(8), 1364-1372. <https://doi.org/10.1093/ps/85.8.1364>
- Hassan, H. M., Samy, A., Youssef, A. W., & Mohamed, M. A. (2018). Research article using different feed additives as alternative to antibiotic growth promoter to improve growth performance and carcass traits of broilers. *International Journal of Poultry Science*, 17, 255-261. <https://doi.org/10.3923/ijps.2018.255.261>
- Hume, M. E., Clemente-Hernández, S. E. R. G. I. O., & Oviedo-Rondón, E. O. (2006). Effects of feed additives and mixed *Eimeria* species infection on intestinal microbial ecology of broilers. *Poultry Science*, 85(12), 2106-2111. <https://doi.org/10.1093/ps/85.12.2106>
- Ibrahim, R. W., Haddad, N. I., Haddadin, M., Salah, K. K. A., & Amri, A. (2012). Diversity among and within wild populations of *Origanum syriacum* collected from Jordan. *Crop Research*, 43(1to3), 249-259.
- Ibrahim, R. W., Haddad, N., Haddadin, M., & Abu-Salah, K. (2012). Evaluation of agro-morphological characters and oil percentage of *Origanum syriacum* L. and *Origanum majorana* L. at three dates of initial cutting. *Jordan Journal of Agricultural Sciences*, 8(1), 33-44.
- Jamroz, D., Orda, J., Kamel, C., Wiliczekiewicz, A., Wiertelcki, T., & Skorupińska, J. (2003). The influence of phytogetic extracts on performance, nutrient digestibility, carcass characteristics, and gut microbial status in broiler chickens. *Journal of Animal and Feed Sciences*, 12(3), 583-596. <https://doi.org/10.22358/jafs/67752/2003>
- Jamroz, D., Wiertelcki, T., Houszka, M., & Kamel, C. (2006). Influence of diet type on the inclusion of plant origin active substances on morphological and histochemical characteristics of the stomach and jejunum walls in chicken. *Journal of Animal Physiology and Animal Nutrition*, 90(5-6), 255-268. <https://doi.org/10.1111/j.1439-0396.2005.00603.x>
- Jang, I. S., Ko, Y. H., Yang, H. Y., Ha, J. S., Kim, J. Y., Kang, S. Y., ... & Lee, C. Y. (2004). Influence of essential oil components on growth performance and the functional activity of the pancreas and small intestine in broiler chickens. *Asian-Australasian Journal of Animal Sciences*, 17(3), 394-400. <https://doi.org/10.5713/ajas.2004.394>
- Janz, J. A. M., Morel, P. C. H., Wilkinson, B. H. P., & Purchas, R. W. (2007). Preliminary investigation of the effects of low-level dietary inclusion of fragrant essential oils and oleoresins on pig performance and pork quality. *Meat Science*, 75(2), 350-355. <https://doi.org/10.1016/j.meatsci.2006.06.027>
- Jaradat, N. A. (2015). Review of the taxonomy, ethnobotany, phytochemistry, phytotherapy and phytotoxicity of germander plant (*Teucrium polium* L.). *medicine*, 3(4).
- Jiang, J., & Xiong, Y. L. (2016). Natural antioxidants as food and feed additives to promote health benefits and quality of meat products: A review. *Meat science*, 120, 107-117. <https://doi.org/10.1016/j.meatsci.2016.04.005>
- Kokkini, S. (1996, May). Taxonomy, diversity and distribution of *Origanum* species. In *Proceedings of the IPGRI international workshop on Oregano* (Vol. 8, p. 12). CIHEAM: Valenzano (Bari), Italy, IPGRI.
- Kosar, M., Dorman, H. J. D., Bachmayer, O., Baser, K. H. C., & Hiltunen, R. (2003). An improved on-line HPLC-DPPH method for the screening of free radical scavenging compounds in water extracts of Lamiaceae plants. *Chemistry of natural Compounds*, 39, 161-166. <https://doi.org/10.1023/A:1024853628326>
- Króliczewska, B., Graczyk, S., Króliczewski, J., Pliszczak-Król, A., Miśta, D., & Zawadzki, W. (2017). Investigation of the immune effects of *Scutellaria baicalensis* on blood leukocytes and selected organs of the chicken's lymphatic system. *Journal of animal science and biotechnology*, 8(1), 1-12. <https://doi.org/10.1186/s40104-017-0152-x>
- Leal, L. N., Beltrán, J. A., Alonso, V., Bello, J. M., den Hartog, L. A., Hendriks, W. H., & Martín-Tereso, J. (2018). Dietary vitamin E dosage and source affects meat quality parameters in light weight lambs. *Journal of the Science of Food and Agriculture*, 98(4), 1606-1614. <https://doi.org/10.1002/jsfa.8635>

- Lee, K. W., Everts, H., Kappert, H. J., Frehner, M., Losa, R., & Beynen, A. C. (2003). Effects of dietary essential oil components on growth performance, digestive enzymes and lipid metabolism in female broiler chickens. *British poultry science*, 44(3), 450-457. <https://doi.org/10.1080/0007166031000085508>
- Li, P., Piao, X., Ru, Y., Han, X., Xue, L., & Zhang, H. (2012). Effects of adding essential oil to the diet of weaned pigs on performance, nutrient utilization, immune response and intestinal health. *Asian-Australasian Journal of Animal Sciences*, 25(11), 1617. <https://doi.org/10.5713%2Fajas.2012.12292>
- Li, W. J., Zhao, G. P., Chen, J. L., Zheng, M. Q., & Wen, J. (2009). Influence of dietary vitamin E supplementation on meat quality traits and gene expression related to lipid metabolism in the Beijing-you chicken. *British Poultry Science*, 50(2), 188-198. <https://doi.org/10.1080/00071660902755409>
- Liu, Q., Lanari, M. C., & Schaefer, D. M. (1995). A review of dietary vitamin E supplementation for improvement of beef quality. *Journal of animal science*, 73(10), 3131-3140. <https://doi.org/10.2527/1995.73103131x>
- Mahmoudi, R., & Nosratpour, S. (2013). Teucrium polium L. essential oil: phytochemical component and antioxidant properties. *International food research Journal*, 20(4).
- Al-Hijazeen, M. A., Al-Rawashdeh, M. S., & Al-Rabadi, G. J. (2022). Cooked broiler meat quality affected by different Mediterranean medicinal plants in the diet. *Animal Bioscience*, 35(2), 290. <https://doi.org/10.5713%2Fab.21.0264>
- Mitsch, P., Zitterl-Eglseer, K., Köhler, B., Gabler, C., Losa, R., & Zimpernik, I. (2004). The effect of two different blends of essential oil components on the proliferation of Clostridium perfringens in the intestines of broiler chickens. *Poultry science*, 83(4), 669-675. <https://doi.org/10.1093/ps/83.4.669>
- Mohammadi Gheisar, M., & Kim, I. H. (2018). Phytobiotics in poultry and swine nutrition—a review. *Italian journal of animal science*, 17(1), 92-99. <https://doi.org/10.1080/1828051X.2017.1350120>
- Gheisar, M. M., Hosseindoust, A., & Kim, I. H. (2015). Evaluating the effect of microencapsulated blends of organic acids and essential oils in broiler chickens diet. *Journal of Applied Poultry Research*, 24(4), 511-519. <https://doi.org/10.3382/japr/pfv063>
- Al-Rabadi, N., Haddad, R., Al-Hijazeen, M. A., Massoh, M., Alqudah, J. M., Jiries, A. G., ... & Al-Rabadi, G. J. (2020). Effect of Garlic powder Supplementation level at different growth stages on Broiler performance. *Bull. Env. Pharmacol. Life Sci*, 9(11), 67-76.
- Oviedo-Rondón, E. O., Hume, M. E., Hernández, C., & Clemente-Hernández, S. (2006). Intestinal microbial ecology of broilers vaccinated and challenged with mixed Eimeria species, and supplemented with essential oil blends. *Poultry science*, 85(5), 854-860. <https://doi.org/10.1093/ps/85.5.854>
- Park, J. H., Kang, S. N., Shin, D., & Shim, K. S. (2015). Antioxidant enzyme activity and meat quality of meat type ducks fed with dried oregano (Origanum vulgare L.) powder. *Asian-Australasian journal of animal sciences*, 28(1), 79. <https://doi.org/10.5713%2Fajas.14.0313>
- Paszkiewicz, M., Budzyńska, A., Różalska, B., & Sadowska, B. (2012). Immunomodulatory role of polyphenols in plants. *Postępy Higieny i Medycyny Doświadczalnej*, 66, 637-646.
- Peng, Q. Y., Li, J. D., Li, Z., Duan, Z. Y., & Wu, Y. P. (2016). Effects of dietary supplementation with oregano essential oil on growth performance, carcass traits and jejunal morphology in broiler chickens. *Animal Feed Science and Technology*, 214, 148-153. <https://doi.org/10.1016/j.anifeedsci.2016.02.010>
- Platel, K., & Srinivasan, K. (2004). Digestive stimulant action of spices: a myth or reality?. *Indian Journal of Medical Research*, 119(5), 167.
- Rahimi, S., Teymori Zadeh, Z., Torshizi, K., Omidbaigi, R., & Rokni, H. (2011). Effect of the three herbal extracts on growth performance, immune system, blood factors and intestinal selected bacterial population in broiler chickens. *Journal of Agricultural Science and Technology*, 13(4), 527-539.
- Ri, C. S., Jiang, X. R., Kim, M. H., Wang, J., Zhang, H. J., Wu, S. G., ... & Qi, G. H. (2017). Effects of dietary oregano powder supplementation on the growth performance, antioxidant status and meat quality of broiler chicks. *Italian Journal of Animal Science*, 16(2), 246-252. <https://doi.org/10.1080/1828051X.2016.1274243>
- Samuel, K. G., Wang, J., Yue, H. Y., Wu, S. G., Zhang, H. J., Duan, Z. Y., & Qi, G. H. (2017). Effects of dietary gallic acid supplementation on performance, antioxidant status, and jejunum intestinal morphology in broiler chicks. *Poultry Science*, 96(8), 2768-2775. <https://doi.org/10.3382/ps/pex091>
- Şenay, S., İslim, P., & Tugay, A. (2019). Supplementation of natural antioxidants to reduced crude protein diets for Japanese quails exposed to heat stress. *Brazilian Journal of Poultry Science*, 21. <https://doi.org/10.1590/1806-9061-2017-0694>
- Skjerm, D. A. (1997). The antimicrobial activity of *Allium sativum*. Ph.D. Thesis, Cardiff University.
- Song, J., Xiao, K., Ke, Y. L., Jiao, L. F., Hu, C. H., Diao, Q. Y., ... & Zou, X. T. (2014). Effect of a probiotic mixture on intestinal microflora, morphology, and barrier integrity of broilers subjected to heat stress. *Poultry science*, 93(3), 581-588. <https://doi.org/10.3382/ps.2013-03455>

- Surai, P. F., Kochish, I. I., Fisinin, V. I., & Kidd, M. T. (2019). Antioxidant defence systems and oxidative stress in poultry biology: An update. *Antioxidants*, 8(7), 235. <https://doi.org/10.3390/antiox8070235>
- Suzuki, N., & Lee, Y. C. (2004). Site-specific N-glycosylation of chicken serum IgG. *Glycobiology*, 14(3), 275-292. <https://doi.org/10.1093/glycob/cwh031>
- Tschirch, H. (2000). Use of natural plant extracts as productive enhancers in modern animal rearing practices. *Zeszyty Naukowe Akademii Rolniczej we Wroclawiu. Konferencje (Poland)*.
- Ultee, A., Bennik, M. H. J., & Moezelaar, R. J. A. E. M. (2002). The phenolic hydroxyl group of carvacrol is essential for action against the food-borne pathogen *Bacillus cereus*. *Applied and environmental microbiology*, 68(4), 1561-1568. <https://doi.org/10.1128/AEM.68.4.1561-1568.2002>
- Viveros, A., Chamorro, S., Pizarro, M., Arija, I., Centeno, C., & Brenes, A. (2011). Effects of dietary polyphenol-rich grape products on intestinal microflora and gut morphology in broiler chicks. *Poultry science*, 90(3), 566-578. <https://doi.org/10.3382/ps.2010-00889>
- Windisch, W., Schedle, K., Plitzner, C., & Kroismayr, A. (2008). Use of phytogetic products as feed additives for swine and poultry. *Journal of animal science*, 86(suppl_14), E140-E148. <https://doi.org/10.2527/jas.2007-0459>
- Yang, J. Y., Zhang, H. J., Wang, J., Wu, S. G., Yue, H. Y., Jiang, X. R., & Qi, G. H. (2017). Effects of dietary grape proanthocyanidins on the growth performance, jejunum morphology and plasma biochemical indices of broiler chicks. *animal*, 11(5), 762-770. <https://doi.org/10.1017/S1751731116002056>
- Zahoor, I., Ghayas, A., and Basheer, A. (2018). Genetics and genomics of susceptibility and immune response to necrotic enteritis in chicken: a review. *Molecular Biology Reports*, 45, 31-37. <https://doi.org/10.1007/s11033-017-4138-8>.
- Zhou, Y., Mao, S., and Zhou, M. (2019). Effect of flavonoid baicalein as a feed additive on the growth performance, immunity, and antioxidant capacity of broiler chickens. *Poultry Science*, 98: 2790-2799. <http://dx.org/10.3382/ps/pez071>.