



Unlocking The Potential Of Hemoglobin Enhancement Agents: A Comprehensive Review

Darshit Kumar^{1*}, Dr. Nishkruti Mehta², Dr. Pragnesh Patani³

^{1*,2,3}Khyati College of Pharmacy, Palodia, Ahmedabad

***Corresponding Author:** Darshit Kumar

*Khyati College of Pharmacy, Palodia, Ahmedabad, Email: darshitprajapat7676@gmail.com

Article History	Abstract
Received: 1 Nov 2023 Revised: 25 Nov 2023 Accepted: 20 Dec 2023	Hemoglobin is an important protein found in red blood cells and is responsible for the two-way respiratory carrier, transporting oxygen from the lungs to the tissues and facilitating the return transport of carbon dioxide in the body. The development of hemoglobin enhancers is an area of medical research interest aimed at improving oxygen delivery and potentially treating various health conditions. This review aims to explore the role of hemoglobin activators, their mechanism of action, current applications, and potential benefits. Mostly beetroot, dates and findla are mainly consumed as food additives for hemoglobin. And to help increase hemoglobin.
CC License CC-BY-NC-SA 4.0	Keywords: Hemoglobin, Haemoglobin enhancer, Beetroot, Dates and Findla.

INTRODUCTION:

Transport oxygen from the environment into cells and transport Carbon dioxide and H⁺ are produced in the opposite direction of metabolism which is essential for the survival of vertebrates. The reversible binding of the ligands to the hemoglobin molecule and the proper allosteric interactions between the binding sites considerably boost the carrying capacity of O₂, CO₂, and H⁺ in blood. All vertebrates, with the noteworthy exception of Antarctic icefishes, contain Hh, which significantly reduces the circulatory demand^[1].

One of the proteins that has been researched the most, hemoglobin, has led to a thorough knowledge of the links between its structure and function. Since its structure and activities are well understood, Hh has earned the moniker "honorary enzyme" and serves as a useful model for research on allosteric interactions in other proteins^[1]. Fish Hbs in particular have been the subject of extensive research on vertebrate Hbs, which has continued to disclose fascinating new facets of cellular and molecular regulatory mechanisms. Since fish Hbs were last examined in this book series, significant advancements have been made^[2]. The goal of this chapter is to provide a summary of what is currently known about the molecular structure, conformational variations, allosteric connections, and the many activities of fish Hb.

Red blood cells contain a metalloprotein called hemoglobin (Hb). Oxygen is circulated throughout the body via red blood cells. Hemoglobin serves as the oxygen transporter in the red blood cells of all vertebrates^[3]. Iron is present in hemoglobin, which makes up 96% of the dry weight of a red blood cell. Hemoglobin is a component of every human body. The hemoglobin level is measured in grams (gm) per deciliter (dL) of whole blood. An adult guy with normal hemoglobin levels ranges from 13.8 to 17.2 g/dL. Adult females (non pregnant) should have 12.1–15.1 g/dL of hemoglobin^[3]. The oxygen equilibrium curve (OEC), which displays Hb's oxygen saturation (SO₂) at various partial pressures of oxygen (pO₂), serves as evidence that Hb's principal job is to transfer oxygen (O₂) from the lung to tissues by cooperatively binding and releasing

oxygen^[4]. The Oxygen affinity for Hb is measured by the partial pressure of oxygen (pO₂) at 50% oxygen saturation (SO₂), which is approximately 26 mmHg for normal adult human Hb (HbA)^[4].

Two classical states' equilibrium have been used to describe the Hb function: the relaxed (R) state (liganded Hb), which has a high affinity for Oxygen, and the tense (T) state (unliganded Hb), which has a low affinity for Oxygen, provide a structural foundation for cooperative effects that make Oxygen absorption and release *in vivo* more efficient^[5,6,7].

STRUCTURE OF HEMOGLOBIN:

Hemoglobin comprises four subunits, each having one globulin chain (polypeptide chain) and one heme group^[8]. Each globulin chain includes an essential iron-containing porphyrin molecule called heme. An iron atom contained within the heme molecule is essential for carrying oxygen and carbon dioxide in human blood. The iron in hemoglobin is also responsible for the red color of blood^[8]. Hemoglobin is made up of four protein molecules (globulin chains) that are linked together; Two alpha-globulin chains. Two beta-globulin chains (Fig. 1)^[8,9]. Similar in length but with different amino acid sequences are the alpha and beta chains. Human hemoglobins, whether they are adult or embryonic, all have the same alpha chain. The gamma chain of fetal hemoglobin ($\alpha_2\gamma_2$), the beta chain of adult-normal hemoglobin ($\alpha_2\beta_2$), and the delta chain of HbA₂ are among the non-alpha chains^[9].




Fig.1: Crystal structure of hemoglobin. a Overall quaternary structure of Hb with the two α chains and β chains colored grey and tan, respectively. b Structure of oxygenated (R state) Hb (magenta) superimposed on the structure of deoxygenated (T state) Hb (blue). Note the larger central water cavity in the T structure.

The two sites of entrance into the central water cavity that are greater in T state structure than R state structure are the α - and β -clefts. More salt-bridge/hydrogen-bond interactions than R state structure are also present at the interdimer interface ($\alpha 1\beta 1$ – $\alpha 2\beta 2$) of the T state structure^[9]. The 7 and 8 helices, referred to respectively as A-H, that make up the α -subunits and β -subunits are connected by non-helical segments. The E and F helices of each subunit provide a binding pocket for heme. The four nitrogen atoms of the porphyrin ring work together to coordinate the ferrous ion stored in the middle to form the heme. an imidazole of a histidine residue found on the F helix (proximal histidine or His (F8)), the Fe is further covalently bound to Hb at the heme proximal pocket^[10].

Through hydrogen-bond interaction, a histidine residue (His E7) in the distal pocket stabilizes the attached Oxygen. A very weakly bound water molecule fills the α -cleft of deoxygenated hemoglobin, leaving a warped octahedron in its place^[10]. central water cavity, Allosterity is a result of cooperativity events that impact the T $\rightarrow$ R transition and are sparked by α - and β -clefts, as well as salt-bridge/hydrogen-bond interactions across the $\alpha 1\beta 1$ – $\alpha 2\beta 2$ ($\alpha 1\beta 2$ or $\alpha 2\beta 1$ or $\alpha 1\alpha 2$ or $\beta 1\beta 2$) dimer interface^[11,12].

Some Fruits help to increase hemoglobin levels:**Beetroot:**

Beta vulgaris, a member of the Chenopodiaceae family and a well-known plant, is commonly known as beetroot^[13,14]. Beetroot is well-known for its therapeutic benefits and usefulness as a juice. In the past, beetroot was used as an aphrodisiac and to increase the hormones involved in sexual activity. To remove kidney and bladder stones, beetroot juice is utilized as a natural therapy^[14,15]. Beetroot has received a lot of attention as a meal that promotes good health^[16,17]. Beetroot is cultivated in several nations across the world, widely consumed as part of the standard diet, and frequently utilized in food manufacture as a food coloring ingredient^[18,19].

Beetroot nutrient include folate – a vitamin that helps keep your blood vessels healthy and Potassium to help protect your Heart^[20].

Beets are also an excellent source of:

- **Folate**
- **Manganese**
- **Vitamin C**
- **Vitamin A**
- **Potassium**

Ascorbic acid, carotenoids, phenolic acids, and flavonoids are among the many phytochemical components found in beetroot^[21,22]. The group of extremely bioactive pigments known as betalains is found in only a few crops, including beetroot. It is classified as either betacyanin pigments, which are red-violet, or other members of the betalain family^[23,24]. Beets contain nitrate in addition to flavonoids, anthocyanins, and betaine, all of which have cardioprotective properties^[25,26]. The plant *B. vulgaris* is a high source of betaine, which has physiological benefits include lowering plasma homocysteine levels, shielding the liver from toxins, and modifying the kidneys' osmolyte function^[27]. Beetroot is a rich and nutritious source, and it has health-promoting properties such as antioxidant and anti-inflammatory properties and hemoglobin boosters^[28]. Beetroots were primarily dietary sources of nitrates, May have important implications for managing cardiovascular health^[29].

Antioxidant:

Recently, antioxidants of the class of betalain pigments were identified. Due to its phenolic components, beetroot has been shown in several published research investigations to possess high antioxidant effects^[30]. betalains and a significant amount of superoxide, hydroxyl anion, and superoxide radical scavenging action towards stable DPPH. Beet root pomace, a byproduct of the juice industry that is inedible yet likely a significant source of antioxidants, is a waste product from beet root harvesting^[31].

Anti-inflammatory activity:

A rat model of paw oedema caused by carageenan was used to test the beta vulgaris plant's anti-inflammatory effects. A 1000 mg/kg dosage of an aqueous extract from the plant's leaves significantly reduced the paw oedema. Beta vulgaris exhibited anti-inflammatory activity in response to foreign body-induced inflammation^[32,33].

Anti-hypertensive:

Heart disease is largely brought on by high blood pressure, a widespread public health issue. According to recent studies, consuming beetroot in bread goods or as a supplement to juice lowers both systolic and diastolic blood pressure. It has been more than 30 years since research on red beet juice's ability to lower blood pressure and enhance the physiological response to exercise was published^[34,35,36].

Dates:

The largest agricultural crop is the date palm (*Phoenix dactylifera*), which is widely distributed across the Arabian Gulf region, including Saudi Arabia, Northern Africa, Pakistan, India, and the United States (California)^[37]. Dates are mostly composed of carbohydrates, which may make about 70% of the total weight (primarily the sugars sucrose, glucose, and fructose). In addition to being a wonderful source of fiber, dates also provide a lot of essential vitamins and minerals, including sizeable levels of calcium, iron, fluorine, and selenium^[38,39,40]. The fruits, bark, and leaves of the date palm are the portions, and they have a wide range of industrial and medical uses. When mature, date fruits have a sugar content ranging from 72 to 88%, making

them a high-energy food source^[41]. Date fruit is rich in phytochemicals such as carotenoids, polyphenols (e.g., phenolic acids, isoflavons, lignans, and flavonoids), tannins, and sterols^[42]. In this way, they are responsible for their antioxidant and antibacterial activities, anti-inflammatory properties, and other health benefits like haemoglobin^[43].




Figure 1. (a) Date palm tree and (b) Date fruit

Carotenoids:

The lipid components of the date fruit include carotenoids, which are regarded as a significant family of phytonutrients. They serve as precursors of vitamin A, which is essential for eyesight and acts as an antioxidant to shield cells from the harmful effects of free radicals^[44]. Dates include lutein, β -carotene, zeaxanthin, and neoxanthin as their main carotenoids. In comparison to red-colored date variations, the yellow color date variety had greater total carotenoids. According to reports, the yellow fruits also contain a complex blend of carotenol fatty acid esters^[45].

Phenolic Acids:

Phenolic acids, which include an aromatic benzene ring with an attached hydroxyl function and one or more carboxylic acid groups, are among the major secondary aromatic plant metabolites. Dates contain a significant amount of phenolic acids, according to several study organizations^[46,47]. Phenolic acids, which are mostly found in the bound form, may be found in a variety of dates in varying amounts and patterns. Fresh dates had an average total phenolic level of 193.7 mg/100 g, whereas dried dates had a total phenolic value of 239.5 mg/100 g^[48].

Phytosterols and Phytoestrogens:

In the lipid-soluble portion of the date fruit, phytosterols are yet another important phytochemical. With chemical structures that are comparable to those of cholesterol, these molecules are only found in plants^[49]. Phytoestrogens are organic substances that have the ability to bind to estrogen receptors and have a variety of estrogenic or antiestrogenic effects^[50].

Findla (Prickly Pears):

Opuntia ficus-indica (L.) Mill., sometimes known as the prickly pear or nopal cactus, is a member of the dicotyledonous angiosperm Cactaceae family. A tropical and subtropical plant is *O. ficus indica*^[51]. Locally it is called as nagphani or danda thohar. It has different names in India like Hathlo thor, chorhthlo (Gujarati), Haththathoira, Nagphana, Nagphani (Hindi), Snuhi, Vajrakantaka, Bahushala (Sanskrit), Nagadali, Nagakkali, Chapati balli (Tamil), Nagamulla, Nagajemudu (Telugu), Nagphani, Thuar (Urdu)^[52]. The nutritional makeup of foods is frequently utilized for nutritional education, assessing the effects of food choices on public health, assessing dietary consumption, and creating and implementing specific recommendations for eating

behaviors^[53]. D-glucose, D-galactose, L-arabinose, D-xylose, L-rhamnose, as well as D-galacturonic and glucuronic acids, are all present in plant mucilage^[54]. Findla has many uses, such as anti-aging, anti-inflammatory, brightening skin tone, and making skin smoother. It is also helpful in healing cuts, wounds, blemishes, and reducing dark circles^[55].



Figure 2. Findla (Prickly pear) fruits.

Conclusion:

Overview of the molecular structure of Hb, relating to its primary function of oxygen transport. Beetroot, dates, and Findla have the ability to improve the carrying of oxygen and carbon dioxide in human blood and also help in the enhancement of hemoglobin. Hemoglobin gives red color to blood. It is provided with folate, vitamins, and minerals. It is a functional food with huge health benefits: antioxidant, anti-inflammatory, anti-hypertensive, and other chronic metabolomics activity. Beetroot, which has been primarily a source of dietary nitrate, may have important implications for managing cardiovascular health. In these cells, hemoglobin performs the dual roles of an antioxidant and a regulator of iron metabolism. Hemoglobin's affinity for oxygen, which can be used to treat disorders brought on by hypoxia and ischemia as well as sickle cell disease.

References:

1. F.B Jensen, Angel fago, Roy E. Weber; Hemoglobin structure and Function. *Fish Respiration*;1998;17: 1-40.
2. Riggs, A., Properties of fish hemoglobins. In "Fish Physiology," Hoar, W. S., and Randall, D. J., eds.1970; 4 : 209-252.
3. Lankna Panawala; What is the Function of Hemoglobin in the Human Body; *researchgate*.2017; 1-7.
4. Mostafa H. Ahmed, Mohini S. Ghatge, Martin K. Safo; Hemoglobin: Structure, Function and Allostery. *Subcell Biochem*.2020 ; 94; 345–382.
5. Perutz MF, Stereochemical mechanism of cooperative effects in haemoglobin. *Biochimie*.1972; 54:587–588.
6. Perutz MF, Wilkinson AJ, Paoli M, Dodson GG, The stereochemical mechanism of the cooperative effects in hemoglobin revisited. *Annu Rev Biophys Biomol Struct*. 1998; 27; 1–34.
7. Safo MK, Ahmed MH, Ghatge MS, Boyiri T. Hemoglobin-ligand binding: understanding Hb function and allostery on atomic level. *Biochim Biophys Acta*. 2011;797–809.
8. Charles Patrick Davis; Hemoglobin: Normal, High, Low Levels, Age & Gender; *Medscape*.2022; 526-531.
9. Alain J. Marengo-Rowe; Structure-function relations of human hemoglobins; *Bayl Univ Med Cent*. 2006;19: 239–245.
10. Winslow RM, Charache S. Hemoglobin Richmond. Subunit dissociation and oxygen equilibrium properties. *J Biol Chem*. 1975; 250; 6939–6942.
11. Baldwin J, Chothia C.;Haemoglobin: The structural changes related to ligand binding and its allosteric mechanism. *J Mol Biol*.1979; 129:175–220.
12. Pagel PS, Hettrick DA, Montgomery MW et al. (1998) RSR13, a synthetic allosteric modifier of hemoglobin, enhances recovery of stunned myocardium in dogs. *Adv Exp Med Biol*. 1998; 454:527–531.

- 13.Dlim, MM, SG Alsabri, SS Mohamed, AE Zetrini and SB Mohamed. Use of Beta vulgaris as natural coloring agent for foods and cosmetics in Libya. *Journal of Chemical and Pharmaceutical Research*.2013; 5(11): 340-345.
- 14.Rajmeet Singh, Jaswinder Kaur, Gurinder Singh; A REVIEW ON BETA VULGARIS. *JOURNAL PUNJAB ACADEMY OF SCIENCES*. 2022; 22(1); 34-40.
- 15.Sharma, N, BS Tanwer, and R Vijayvergia. Study of medicinal plants in Aravali regions of Rajasthan for treatment of kidney stone and urinary tract troubles. *International Journal of PharmTech Research* 2011; 3(1):110–113.
- 16.Ninfali, P.; Angelino, D. Nutritional and functional potential of Beta vulgaris cicla and rubra. *Fitoterapia* 2013; 89: 188–199.
- 17.Tom Clifford,Glyn Howatson,Daniel J. West and Emma J. Stevenson; The Potential Benefits of Red Beetroot Supplementation in Health and Disease; *Open access nutriens* 2015; 7: 2801-2822.
- 18.Georgiev, V.G.; Weber, J.; Kneschke, E.M.; Denev, P.N.; Bley, T.; Pavlov, A.I. Antioxidant activity and phenolic content of betalain extracts from intact plants and hairy root cultures of the red beetroot Beta vulgaris cv. Detroit dark red. *Plant Foods Hum. Nutr.* 2010; 65: 105–111.
- 19.Zielińska-Przyjemská, M.; Olejnik, A.; Dobrowolska-Zachwieja, A.; Grajek, W. In vitro effects of beetroot juice and chips on oxidative metabolism and apoptosis in neutrophils from obese individuals. *Phytosphaera Res.* 2009; 23: 49–55.
- 20.Christine Mikstas; Health Benefits of Beetroot;2022. Beetroot: Health Benefits, Nutrients, Preparation, and More (webmd.com)
- 21.Kujala, T.S.; Vienola, M.S.; Klika, K.D.; Loponen, J.M.; Pihlaja, K. Betalain and phenolic compositions of four beetroot (Beta vulgaris) cultivars. *Eur. Food. Res. Technol.* 2002, 214, 505–510.
- 22.Wootton-Beard, P.C.; Ryan, L. A beetroot juice shot is a significant and convenient source of bioaccessible antioxidants. *J. Funct. Foods* 2011, 3, 329–334.
- 23.Lee, C.H.; Wettasinghe, M.; Bolling, B.W.; Ji, L.L.; Parkin, K.L. Betalains, phase II enzyme-inducing components from red beetroot (Beta vulgaris L.) extracts. *Nutr. Cancer* 2005, 53, 91–103.
- 24.Vulic, J.J.; Cebovic, T.N.; Canadanovic-Brunet, J.M.; Cetkovic, G.S.; Canadanovic, V.M.; Djilas, S.M.; Tumbas Saponjac, V.T. In vivo and in vitro antioxidant effects of beetroot pomace extracts. *J. Funct. Foods*. 2014, 6, 168–175.
- 25.Zhao G, He F, Wu C, Li P, Li N, Deng J, Zhu G, Ren W, Peng Y. Betaine in inflammation: Mechanistic aspects and applications. *Front Immunol.* 2018 ;9 :1070.
- 26.Cassidy A, Mukamal KJ, Liu L, Franz M, Eliassen AH, Rimm EB. High anthocyanin intake is associated with a reduced risk of myocardial infarction in young and middle-aged women. *Circulation*. 2013;127(2):188–96.
- 27.Szafer, H., Krajka-Kuźniak, V., Ignatowicz, E., Adamska, T., Baer-Dubowska, W; Evaluation of the effect of beetroot juice on DMBA-induced damage in liver and mammary gland of female Sprague–Dawley rats. *Phytother. Res.*2014; 28, 55–61.
- 28.Georgiev VG, Weber J, Kneschke E-M, Denev PN, Bley T, Pavlov AI. Antioxidant activity and phenolic content of Betalain extracts from intact plants and hairy root cultures of the red beetroot Beta vulgaris cv. Detroit dark red. *Plant Foods Hum Nutr.* 2010;65(2):105–11.
- 29.Lundberg, J.O.; Weitzberg, E.; Gladwin, M.T. The nitrate-nitrite-nitric oxide pathway in physiology and therapeutics. *Nat. Rev.* 2008, 7, 156–167.
- 30.Kapur, A.N., Sati, S.A., Ranjan, A.S., Gupta, P.R; Screening methanolic extracts of beta vulgaris roots for photoprotective activity. *Int. J. Pharm. Pharm. Sci.* 2012; 4, 124–127.
- 31.CB, JM, SS, GS and VJJ; Antioxidant and antimicrobial activities of beet root extracts. *Czech J. Food Science*.2011; 29: 575– 585.
- 32.Marjan Nikan and azadeh Manayi ; Beta Vulgaris L. *Medical Plant Research Center* ; 2019;153-157.
- 33.Jain, SGV and PK Sharma. Anti-inflammatory activity of aqueous extract of Beta vulgaris L. *Journal of Basic and Clinical Pharmacy*. 2011; 2: 83–86.
- 34.Hobbs, D.A., George, T.W., Lovegrove, J.A.; The effects of dietary nitrate on blood pressure and endothelial function: a review of human intervention studies. *Nutr. Res. Rev.*2013a; 26, 210–222.
- 35.Jajja, A., Sutyarjoko, A., Lara, J., Rennie, K., Brandt, K., Qadir, O., Siervo, M; Beetroot supplementation lowers daily systolic blood pressure in older, overweight subjects. *Nutr. Res.*2014; 34, 868–875.
- 36.Wylie, L.J., Kelly, J., Bailey, S.J., Blackwell, J.R., Skiba, P.F., Winyard, P.G., Jeukendrup, A.E., Vanhatalo, A. and Jones, A.M.; Beetroot Juice and Exercise: Pharmacodynamic and Dose-Response Relationships. *Journal of Applied Physiology*.2013; 115, 325-336.

37. W ghori , N Saba , M Jawaaid and M Asim ; A review on date Palm(Phoenix dactylifera) fibers and its Polymer Composites. *Open Access*; 2018; 1-17.
38. Al-Farsi, M. and Lee, C.Y.; Nutritional and functional properties of dates: A review. *Critical Review in Food Science and Nutrition*. 2008; 48: 877–884.
39. Al-Shahib, W. and Mvarshall, R.J.; The fruit of date palm: its possible uses as best food for the future. *International Journal of Foods Science and Nutrition*.;2003;354:247–259.
40. Khan, M., Sarwar, A., Wahab, M., and Haleem, R.; Physio-chemical characterization of date varieties using multivariate analysis. *Journal of Food Agriculture*.;2008; 88: 1051–1059.
41. Ahmed I.A., Ahmed A.W.K., Robinson R.K.; Chemical composition of date varieties as influenced by the stage of ripening. *Food Chem*.;1995; 54, 305–309.
42. Martín-Sánchez, A. M., Cherif, S., Ben-Abda, J., Barber-Vallés, X., PérezÁlvarez, J. Á., and Sayas-Barberá, E.; Phytochemicals in date co-products and their antioxidant activity. *Food Chem*.;2014; 158, 513–520.
43. El-Far, A. H., Oyinloye, B. E., Sepehrimanesh, M., Allah, M. A. G., Abu-Reidah, I., Shaheen, H. M., et al.; Date palm (Phoenix dactylifera): Novel findings and future directions for food and drug discovery. *Current Drug Discovery Technologies*;2019; 16 (1), 2–10.
44. Julia, V., Macia, L., and Dombrowicz, D.; The impact of diet on asthma and allergic diseases. *Nat. Rev. Immunol*.;2015; 15, 308–322.
45. Amanat Ali, Mostafa Waly, M. Mohamed Essa, and Sankar Devaraja; Nutritional and Medicinal Value of Date Fruit; *researchgate*; 2012;361-375.
46. Benmeddour, Z., Mehinagic, E., Le Meurlay, D., and Louaileche, H.; Phenolic composition and antioxidant capacities of ten Algerian date (Phoenix dactylifera L.) cultivars: a comparative study. *J. Funct. Food* ; 2013; 346–354.
47. El Sohaimy, S., Abdelwahab, A., Brennan, C., and Aboul-enein, A.; Phenolic content, antioxidant and antimicrobial activities of Egyptian date palm (Phoenix dactylifera L.) fruits. *Aust. J. Basic Appl. Sci.* 9; 2015; 141–147.
48. Al-Farsi, M., Alasalvar, C., Morris, A., Baron, M., and Shahidi, F.; Comparison of antioxidant activity, anthocyanins, carotenoids, and phenolic of three native fresh and sun dried date (Phoenix dactylifera L.) varieties grown in Oman. *Journal of Agriculture and Food Chemistry*; 2005b; 53: 7592–7599.
49. Al-Laith, A. A.; Degradation kinetics of the antioxidant activity in date palm (Phoenix dactylifera L.) fruit as affected by maturity stages. *Arab Gulf J. Sci. Res*.;2009; 27: 16–25.
50. Al-Turki, S., Shahba, M. A., and Stushnoff, C. J.; Diversity of antioxidant properties and phenolic content of date palm (Phoenix dactylifera L.) fruits as affected by cultivar and location. *J. Food Agri. Environ*.; 2010; 8: 253–260.
51. Butera, D.; Tesoriere, L.; di Gaudio, F.; Bongiorno, A.; Allegra, M.; Pintauro, A.M.; Kohen, R.; Livrea, M.A. Antioxidant activities of sicilian prickly pear (Opuntia ficus indica) fruit extracts and reducing properties of its betalains: Betanin and indicaxanthin. *J. Agric. Food Chem.* 2002; 50; 6895–6901.
52. Chauhan SP, Sheth NR, Jivani NP, Rathod IS, Shah PI. Biological actions of Opuntia species. *Syst Rev Pharm.* 2011; 1(2): 146- 151.
53. Costa, H.S., et al.; New nutritional data on traditional foods for European food composition databases. *Eur. J. Clin. Nutr*.;2010; 64 (S3), S73-S81.
54. El-Samahy S K, El-Hady EAA, Habiba RA, and Moussa TE. Chemical and Rheological Characteristics of Orange-Yellow Cactus-Pear Pulp from Egypt. *J. PACD.* 2006; 39-51.
55. Alexandra Dusenberry; 31 Benefits Of Prickly Pear Fruit For Skin, Hair & Health; stylecraze; 2023. <https://www.stylecraze.com/articles/benefits-of-prickly-pear-for-skin-hair-and-health>.