

Ecological Interactions of Invasive Insects and Native Plant Species in Changing Climate

Manish K Yadav¹, Satya Narayan Satapathy^{2*}, Chandan Kumar Panigrahi³, Pritam Das⁴, Pritendra Narayan Paikaray⁵, Simran Mahapatra⁶, Samit Pal⁷, and Subhashree Subhasmita Paikaray⁸

¹Assistant Professor, Entomology MS Swaminathan School of Agriculture, CUTM Paralakhemundi Odisha 761211

^{2*}Assistant Professor, Department of Entomology Faculty of Agricultural Sciences, Siksha 'O' Anusandhan Deemed to be University, Bhubaneswar, Odisha, India.

³PG Scholar, Entomology, SHUATS, Prayagraj, UP, India.

^{4,5}Teaching Assistant, Department of Pathology Faculty of Agricultural Sciences, Siksha 'O' Anusandhan Deemed to be University, Bhubaneswar, Odisha, India.

^{6&7}PG Scholar, Department of Entomology Faculty of Agricultural Sciences, Siksha 'O' Anusandhan Deemed to be University, Bhubaneswar, Odisha, India.

⁸PhD Scholar, Department of Entomology Faculty of Agricultural Sciences, Siksha 'O' Anusandhan Deemed to be University, Bhubaneswar, Odisha, India.

Email: satyanarayansatapathy40@gmail.com¹

*Corresponding author's E-mail: satyanarayansatapathy40@gmail.com

Article History	Abstract
Received: 15 June 2023 Revised: 10 Sept 2023 Accepted: 10 Oct 2023	<i>Invasive insects pose a significant threat to native plant species and ecosystems, particularly in the context of changing climates. Understanding these interactions is crucial for effective conservation and management strategies aimed at mitigating the adverse effects of invasive species on native plant communities. Invasive insects often establish and proliferate in new habitats due to the absence of natural enemies and the availability of suitable resources. As climate change alters the distribution and phenology of plants, it can influence the susceptibility and resilience of native plant species to invasive insects. In some cases, rising temperatures and altered precipitation patterns may favour the spread and population growth of invasive insects, leading to increased herbivory, reduced plant fitness, and ultimately, altered community dynamics. Furthermore, changing climates can disrupt the synchrony between native plants and their pollinators or beneficial insect populations, further exacerbating the impacts of invasive insects. As native plants and pollinators respond differently to shifting climatic conditions, their interactions may become disrupted, potentially reducing the reproductive success and long-term survival of native plant populations. However, it is important to note that climate change can also create novel opportunities for both invasive insects and native plant species. In certain instances, invasive insects may benefit from warmer temperatures and expanded ranges, while some native plants may exhibit adaptive responses and resilience to changing climatic conditions. These complex interactions highlight the need for a comprehensive understanding of the ecological dynamics between invasive insects and native plant species under various climate scenarios. The ecological interactions between invasive insects and native plant species in changing climates have far-reaching consequences for biodiversity conservation and ecosystem functioning. As climates continue to evolve, it is imperative to further investigate these interactions and develop adaptive strategies to mitigate the impacts of invasive insects on native plant communities. By doing so, we can strive to preserve and restore ecological balance in the face of ongoing environmental change.</i>
CC License CC-BY-NC-SA 4.0	Keywords: introduced pests, invasion, quarantine, endemic, pests, behaviour, insects

1. Introduction

Invasive insect pests pose significant threats to agricultural systems worldwide, leading to substantial economic losses and ecological disruptions. Understanding the dynamics between invasive insects and agricultural practices is essential for developing effective strategies to mitigate their negative effects and promote sustainable food production (Tonnang *et al.*, 2022). Invasive insect pests have the ability to rapidly establish and proliferate in new regions, often surpassing the natural defences of local ecosystems. They can inflict direct damage to crops through feeding, oviposition, and transmission of pathogens, causing yield losses and reduced quality of agricultural products. Furthermore, these pests can disrupt the balance of agroecosystems by altering nutrient cycling, plant-pollinator interactions, and natural pest control mechanisms (Gupta *et al.*, 2019; Fortuna *et al.*, 2022). Climate change, globalization, and increased international trade have facilitated the introduction and spread of invasive insect pests, intensifying the challenges faced by agricultural systems. Rising temperatures, altered precipitation patterns, and shifting phenological events can create more favourable conditions for invasive species, enabling them to expand their ranges and infest new areas (Mallet, 2018). Additionally, the movement of goods and people across borders increases the likelihood of pest introductions, highlighting the importance of strict biosecurity measures and regulations. Managing invasive insect pests in agriculture requires a multifaceted approach that integrates various strategies (Skendžić *et al.*, 2021). Integrated Pest Management (IPM) practices, which combine cultural, biological, and chemical control methods, are commonly employed to minimize the use of pesticides and promote environmentally sustainable pest management. Monitoring and early detection systems play a crucial role in identifying invasive species at their early stages of establishment, enabling timely interventions and preventing their widespread impacts (Musolin *et al.*, 2022).

Biological control, involving the introduction or augmentation of natural enemies, has proven effective in managing some invasive insect pests. However, careful consideration must be given to the potential ecological risks associated with the introduction of non-native species for control purposes (Wright, 2014). Additionally, research and innovation in breeding pest-resistant crop varieties, developing pheromone-based trapping systems, and exploring alternative pest control methods, such as the use of microbial agents or precision farming technologies, hold promise for sustainable pest management in agriculture (Venette and Hutchison, 2021). International cooperation and collaboration among countries are vital for effectively addressing the challenges posed by invasive insect pests. Sharing knowledge, best practices, and resources can enhance surveillance and response systems, facilitate the development of early warning systems, and promote coordinated pest management efforts. Furthermore, public awareness and education programs can play a critical role in engaging farmers, stakeholders, and the general public in recognizing and reporting invasive species, thereby contributing to their effective control and prevention (Gupta *et al.*, 2019). Invasive insect pests pose significant threats to agriculture, necessitating comprehensive and integrated approaches to their management. By implementing sustainable pest management practices, strengthening biosecurity measures, and promoting international cooperation, it is possible to mitigate the impacts of invasive species on agricultural systems and safeguard food production. Continued research and innovation are crucial for developing new strategies and technologies to effectively manage invasive insect pests, ensuring the resilience and sustainability of global agriculture in the face of ongoing challenges (Rathee and Dalal, 2018).

Invasive pests

Invasive pests pose a significant threat to native vegetation and insects, leading to profound ecological disruptions and biodiversity loss. Invasive pests can have detrimental effects on native vegetation by directly damaging plants through feeding, competition for resources, or transmission of pathogens (Daniel *et al.*, 2020). They often lack natural predators or parasites in their new habitats, allowing them to proliferate rapidly and exert considerable pressure on native plant populations. This can result in reduced plant fitness, altered community composition, and, in extreme cases, even the displacement or extinction of native plant species (Bhavani *et al.*, 2019). Furthermore, invasive pests can disrupt the intricate interactions between plants and insects, which are essential for ecosystem functioning. Native insects, including pollinators and herbivores, have evolved in association with native plant species and depend on them for food and habitat. When invasive pests negatively impact native plants, it can have cascading effects on insect populations, leading to declines in pollination services, shifts in herbivory patterns, and altered trophic interactions within ecosystems (Bisht and Giri, 2019). The impacts of invasive pests on native insects are particularly concerning due to the critical roles insects play in ecological processes such as pollination, nutrient cycling, and decomposition. Disruptions to insect populations can have far-reaching consequences for ecosystem health and stability. Additionally, as

native insects are often specialized in their interactions with specific plant species, the loss or decline of native plants can result in the loss of host plant resources and subsequent declines in insect biodiversity (Mani, 2017). Climate change can exacerbate the impacts of invasive pests on native vegetation and insects. Rising temperatures, altered precipitation patterns, and shifting phenology can create more favourable conditions for invasive species, allowing them to expand their ranges and intensify their impacts. Moreover, climate change can disrupt the synchrony between native plants and their associated insects, leading to phenological mismatches and reduced reproductive success for both plants and insects (Tonnang *et al.*, 2022).

To effectively address the impacts of invasive pests on native vegetation and insects, integrated management approaches are crucial. These approaches should include early detection and rapid response systems, targeted pest control methods, restoration efforts for native plant communities, and the promotion of habitat connectivity and biodiversity conservation (Viraktamath, 2002). Additionally, collaboration between scientists, land managers, policymakers, and the public is essential for implementing effective invasive species management strategies. Invasive pests pose significant threats to native vegetation and insects, jeopardizing ecosystem functioning and biodiversity conservation (Kallethwaraswamy, 2018). The impacts of invasive pests on native plant communities and insect populations highlight the urgent need for proactive and coordinated management efforts. By understanding the ecological dynamics between invasive pests, native vegetation, and insects, we can develop strategies to mitigate their negative effects and promote the long-term sustainability of native ecosystems (Singh *et al.*, 2020).

Interaction of native and invasive insects

The interaction between native and invasive insects in ecosystems is a dynamic process that can have profound ecological implications. Native and invasive insects often occupy similar ecological niches, competing for limited resources such as food, habitat, and mates. Competition between native and invasive insects can lead to changes in population dynamics, distribution patterns, and community composition (Rodriguez, 2006). In some cases, invasive insects may outcompete native species, leading to declines in native insect populations and potential disruptions to ecosystem functioning. On the other hand, facilitation can also occur between native and invasive insects. In certain instances, native insects may benefit from the presence of invasive species by utilizing novel resources or habitats made available by the invaders. This facilitation can have both positive and negative consequences, depending on the specific context and the ecological traits of the interacting species (Johnson, 2009).

Predation is another important interaction between native and invasive insects. Native predators may recognize and prey upon invasive insects, potentially exerting some level of control over their populations. However, the effectiveness of native predators in regulating invasive species can be influenced by various factors, including prey preferences, predator-prey interactions, and the ability of invasive insects to defend themselves or escape predation (Jackson, 2015). Indirect effects can also arise from the interaction of native and invasive insects. For example, the introduction of an invasive insect may disrupt the mutualistic relationships between native insects and plants, leading to cascading effects on plant-pollinator interactions or herbivory dynamics. Additionally, the presence of invasive insects can alter the behaviour, physiology, or life history traits of native insects, potentially affecting their survival, reproduction, and overall fitness (Mills *et al.*, 2004).

Understanding the interaction between native and invasive insects is crucial for developing effective pest management strategies. Integrated Pest Management (IPM) approaches, which take into account the ecological context and consider a combination of strategies, can help mitigate the negative impacts of invasive insects while minimizing harm to native species (Bauer, 2012). These strategies may include biological control using native predators or parasitoids, habitat management to enhance native insect populations, and targeted interventions to control invasive species. Furthermore, the conservation of native insect populations and their habitats is essential for maintaining the resilience and biodiversity of ecosystems. Protecting and restoring native plant communities, promoting habitat connectivity, and reducing the use of broad-spectrum insecticides can help support native insect populations, enabling them to withstand the pressures exerted by invasive species (With, 2002). The interaction between native and invasive insects is a complex process with far-reaching ecological implications. Understanding the dynamics of these interactions is crucial for effective pest management, biodiversity conservation, and the maintenance of ecosystem balance. By considering the ecological context and implementing integrated strategies, it is possible to mitigate the negative impacts of invasive insects while preserving native insect populations and the ecological services they provide (Didham, 2005).

Invasive insects are a significant concern for ecosystems, agriculture, and public health. These insects, often introduced to new environments unintentionally or intentionally, can have detrimental effects on the native flora and fauna. While they possess certain abilities that aid their survival and proliferation, it is crucial to recognize the negative impacts they can have and the importance of managing their populations effectively (Rodriguez, 2006). One of the key abilities of invasive insects is their capacity to adapt to new environments quickly. They possess mechanisms that enable them to establish and thrive in habitats that may be different from their native range. These insects can exploit available resources, such as food, shelter, and breeding sites, with great efficiency, allowing them to outcompete native species. Their ability to reproduce rapidly also contributes to their success in colonizing new areas (Johnson *et. al.*, 2009).

Invasive insects can disrupt ecological balance by preying on or outcompeting native species for resources. They can negatively affect biodiversity by driving native species to extinction or altering natural ecosystems. For example, invasive insects like the emerald ash borer in North America have devastated ash tree populations, resulting in the loss of habitat for countless organisms that depend on these trees (Jackson, 2015). The impacts can extend to higher trophic levels, affecting entire food webs. Furthermore, invasive insects often bring with them novel pests and diseases that can affect agricultural production. They can damage crops, reduce yields, and cause economic losses for farmers. Examples include the Asian citrus psyllid, which transmits a bacterium responsible for citrus greening disease, and the brown marmorated stink bug, which damages a wide range of fruits and vegetables. Controlling these pests is essential to safeguarding agricultural productivity and food security (Mills *et. al.*, 2004).

The management of invasive insects is of utmost importance. Early detection and rapid response are crucial to prevent their establishment and spread. Effective strategies may include quarantine measures, surveillance programs, and the implementation of biological control methods that target specific pests while minimizing harm to non-target species (Bauer, 2012). Public awareness and education play a significant role in preventing the unintentional introduction and spread of invasive insects. People should be educated about the risks associated with the transport of potentially invasive species through international trade, travel, or the movement of goods (Crowl *et. al.*, 2008). Furthermore, understanding and recognizing the signs of invasive insect infestations can facilitate early reporting and prompt action. While invasive insects possess certain abilities that contribute to their success, their impact on ecosystems, agriculture, and public health is significant (With, 2002). The ability to adapt to new environments quickly and exploit available resources allows them to establish and thrive, but it also poses a threat to native species and ecosystems. Recognizing the importance of managing invasive insects effectively through early detection, response, and education is crucial to mitigate their negative effects and preserve the health and balance of our ecosystems (Zavaleta *et. al.*, 2001).

Effect of invasive insects on native insect species

Invasive insects can have a profound impact on native insect populations; often disturbing their ecological balance and causing significant disruptions within ecosystems (Hickman *et. al.*, 2013). Here are a few examples that illustrate how invasive insects can negatively affect native insects:

European Honeybees vs. Native Bees

The introduction of European honeybees (*Apis mellifera*) in various regions has had unintended consequences for native bee populations. European honeybees are highly competitive for floral resources and can outcompete native bees for nectar and pollen (Cane, 2003). This competition can lead to a reduction in food availability for native bees, potentially impacting their survival and reproduction (Chantawannakul *et. al.*, 2018). The introduction of *Apis mellifera*, or the European honeybee, has had both positive and negative impacts on native bee populations. While honeybees are highly valued for their pollination services and honey production, their presence can also have unintended consequences for native bees (Kumar & Kumar, 2014). *Apis mellifera* colonies can be large and highly efficient in collecting nectar and pollen from flowers. This can lead to increased competition for floral resources with native bee species. Honeybees have been observed to outcompete native bees for limited resources, potentially reducing the availability of food sources for native bee populations (Kojima *et. al.*, 2011). As a result, native bees may experience decreased foraging success, reduced reproductive rates, and increased vulnerability to other stressors. The abundance and foraging behaviour of honeybees can influence the dynamics of plant-pollinator relationships (Goulson, 2003). Native bees have evolved alongside native plants and have specialized adaptations that allow them to effectively pollinate specific plant species. However, the introduction of honeybees can disrupt these relationships by monopolizing floral resources and reducing the opportunities for native bees to fulfil their ecological roles as

pollinators. This can have cascading effects on plant populations, especially those that rely on specialized native bee pollinators for reproduction (Paini, 2004).

Honeybees can act as carriers and vectors of diseases and parasites that can affect both honeybees and native bees. For instance, the Varroa mite (*Varroa destructor*), a parasitic mite that feeds on honeybee brood, can also infest and negatively impact native bee populations. The spread of pathogens and parasites by honeybees can lead to increased disease pressure on native bee species, potentially causing declines in their populations and compromising their health and survival (Chakrabarti *et al.*, 2015). Hybridization between honeybees and native bee species can occur in areas where the two populations overlap. This can result in genetic introgression, where genes from the introduced honeybees are incorporated into the gene pool of native bees. Hybridization can lead to the loss of unique genetic traits and adaptations in native bee populations, potentially reducing their ability to cope with environmental changes or increasing their susceptibility to certain stressors. It is important to note that the impacts of *Apis mellifera* on native bees can vary depending on local environmental conditions, bee species, and the intensity of honeybee management. Additionally, the effects can be more pronounced in areas where native bee populations are already under pressure due to habitat loss, pesticide exposure, and other factors (Hurn, 1997).

Red Imported Fire Ants vs. Ground-Nesting Insects

Red imported fire ants (*Solenopsis invicta*) are notorious for their aggressive behavior and ability to displace native ground-nesting insects. These invasive ants construct large underground colonies and have been observed to prey upon or outcompete native insects, including ground-nesting bees, wasps, and other arthropods (Vinson, 1997; Drees, 1998; Wojcik *et al.*, 2001). This disruption in the native insect community can have cascading effects on other organisms that rely on these insects for pollination or as a food source (Klotz *et al.*, 2003; Jemal & Jones, 1993; LeBrun *et al.*, 2012).

Asian Longhorned Beetle vs. Native Wood-Boring Insects

The Asian longhorned beetle (*Anoplophora glabripennis*) is a destructive invasive species that attacks a wide range of hardwood trees. As larvae, these beetles bore into the wood, causing extensive damage and potentially killing the host trees (Duan *et al.*, 2016). In regions where the Asian longhorned beetle has become established, it can outcompete and displace native wood-boring insects that depend on the same tree species for their life cycles. This disruption can lead to declines in native wood-boring insect populations, affecting nutrient cycling and decomposition processes in forests (Hu *et al.*, 2009; Marchioro and Faccoli, 2021).

Invasive pest- *Spodoptera frugiperda*

The invasive pest *Spodoptera frugiperda*, commonly known as the fall armyworm, has had significant impacts not only on crops but also on other polyphagous pests within its invaded range. Here are some of the key effects of fall armyworm on other pests:

Competition for Resources

Fall armyworm larvae are voracious feeders and can consume a wide range of host plants, including major cereal crops such as maize, rice, and sorghum. Their presence can lead to increased competition for food resources among other polyphagous pests that feed on similar crops. This competition may result in reduced food availability and nutritional stress for these pests, potentially impacting their development, survival, and population dynamics (Sokame *et al.*, 2020; Divya *et al.*, 2021).

Displacement of Native Pests

In invaded areas, fall armyworm has the potential to outcompete and displace native polyphagous pests that share similar ecological niches. Native pests may face challenges in the presence of fall armyworm due to its aggressive feeding behavior and reproductive capacity. As a result, native pest populations may decline or become less dominant in agricultural landscapes, leading to shifts in pest dynamics and potential changes in pest management strategies (Hailu *et al.*, 2021; Mutyambai *et al.*, 2022).

Altered Pest Management Practices

The invasion of fall armyworm has prompted changes in pest management strategies, including increased pesticide applications and the adoption of new control methods. These changes can have indirect effects on other polyphagous pests. For example, the increased use of broad-spectrum insecticides to control fall armyworm may also impact non-target pests, including beneficial insects that contribute to natural pest control. Consequently, the altered pest management practices may disrupt the overall pest ecology and potentially lead to unintended consequences (Chimweta *et al.*, 2020; Ahissou *et al.*, 2021). Not only in case of insects, even in

other animals like Brown Treesnake vs. Native Birds: The introduction of the brown treesnake (*Boiga irregularis*) to the island of Guam has had devastating consequences for native bird populations (Wiles *et. al.*, 2003). The snakes, which are native to Australia and Indonesia, prey upon bird eggs and nestlings, causing severe declines in native bird species. The loss of native birds has had cascading effects on the ecosystem, including an increase in the population of invasive insects that were previously controlled by the birds (Rodda *et. al.*, 1999). This example highlights how the disruption of native insectivorous species can indirectly impact native insects.

These examples demonstrate the ways in which invasive insects can disturb native insect populations. Whether through direct competition, predation, resource depletion, or habitat modification, invasive insects can significantly alter the dynamics and composition of native insect communities. It is important to understand these impacts to develop effective strategies for managing invasive species and mitigating their effects on native ecosystems. The impact of invasive insects on native plant species is a significant concern for biodiversity conservation and ecosystem stability. Invasive insects, often introduced unintentionally or intentionally to new environments, can have detrimental effects on native plant populations. Invasive insects are often highly efficient herbivores, capable of consuming and damaging native plant species. They may possess specialized feeding adaptations or lack natural predators, allowing them to rapidly proliferate and cause extensive damage. This can result in reduced plant growth, decreased seed production, and even plant mortality (Avanesyan *et. al.*, 2021). Invasive insects can directly consume foliage, stems, flowers, fruits, and seeds, leading to the loss of reproductive potential and altering the structure and composition of native plant communities. Invasive insects can outcompete native herbivores for limited resources, including host plants. Their aggressive feeding habits and ability to reproduce quickly can give them a competitive advantage, leading to a reduction in food availability for native herbivorous insects (Crowder and Snyder, 2010). This can disrupt ecological interactions and potentially cause declines in native herbivore populations, altering the balance of herbivory within ecosystems.

Invasive insects that act as pollinators may compete with native pollinators for floral resources, potentially affecting the reproductive success of native plant species. Invasive pollinators may have different preferences for flower morphology, nectar production, or pollen transfer mechanisms, leading to shifts in plant-pollinator interactions. This can result in reduced pollination services for native plants, leading to decreased seed set and genetic diversity, and potentially hindering the regeneration of native plant populations (Gurevitch and Padilla, 2004). Invasive insects can serve as vectors for pathogens and diseases that can affect native plant species. These insects may carry and transmit pathogens directly to plants while feeding or indirectly through their excretions. Pathogens can cause diseases that weaken or kill native plants, further compromising their ability to compete with invasive species. Invasive insects can introduce novel pathogens to which native plants may have little or no resistance, leading to increased disease prevalence and potential declines in native plant populations (Cushman *et. al.*, 2011).

Native plants play a vital role in ecosystem functioning, providing habitat, food, and shelter for a diverse array of organisms. When invasive insects negatively impact native plant populations, it can have cascading effects on other trophic levels (Lonighi, 2023). Reduced plant biomass and altered plant community composition can affect the abundance and distribution of other organisms, including herbivores, predators, and decomposers (O'Neil *et. al.*, 2023). This disruption of ecosystem functions can lead to changes in nutrient cycling, energy flow, and overall ecosystem resilience. Mitigating the impact of invasive insects on native plant species requires comprehensive management strategies. These may include early detection and rapid response efforts, the development and implementation of biological control measures, promoting the use of native plant species in restoration and landscaping practices, and minimizing the unintentional introduction and spread of invasive insects through international trade and travel. Protecting and conserving native plant species is crucial for maintaining ecosystem health, biodiversity, and the services they provide (Alfonzetti *et. al.*, 2023; Peña *et. al.*, 2023).

Invasive insects and native plant species are key components of ecosystems worldwide. However, with the onset of climate change, their ecological interactions are being altered, posing significant challenges for biodiversity conservation and ecosystem functioning. Invasive insects can have detrimental effects on native plant species, and the changing climate further exacerbates these impacts. This essay aims to explore the ecological interactions between invasive insects and native plant species in the context of a changing climate,

highlighting the potential consequences and implications for ecosystem dynamics and conservation efforts (Chavre and Patil,2023).

Climate Change and Altered Phenology

Climate change influences the phenology, or timing of biological events, of both invasive insects and native plants. Rising temperatures and shifting seasonal patterns can lead to phenological mismatches, where the emergence and activity of invasive insects do not align with the flowering and growth cycles of native plants. This can disrupt essential pollination processes and reduce seed production, affecting the reproductive success and fitness of native plant species (Elias,2023).

Range Expansion and Colonization

Climate change can create favourable conditions for invasive insects, allowing them to expand their ranges into new areas. As invasive insects colonize new regions, they may encounter native plant species that lack natural defences against their feeding or reproductive strategies. The absence of co-evolutionary relationships between invasive insects and native plants can exacerbate the impacts, leading to increased herbivory, reduced plant growth, and potential declines in native plant populations (Sarkar and Chakroborty,2023).

Altered Plant-Insect Interactions

Invasive insects can directly influence the dynamics of native plant communities through herbivory, altering plant growth, reproduction, and overall ecosystem structure. They may also introduce novel pathogens or interact with existing plant diseases, further compromising the health and resilience of native plant species. These disruptions can have cascading effects on other trophic levels, affecting pollinators, herbivores, and predators that depend on native plants for their survival and reproduction (Cushman *et. al.*, 2011).

Feedback Loops and Ecosystem Resilience

The ecological interactions between invasive insects and native plant species, combined with climate change impacts, can create feedback loops that further exacerbate ecosystem disturbances. For example, increased herbivory by invasive insects may weaken native plant populations, making them more susceptible to other stressors such as drought, diseases, or invasive plants. These feedback loops can lead to declines in native plant diversity, compromised ecosystem resilience, and increased vulnerability to further environmental changes (Huet. *al.*,2009; Marchioro and Faccoli,2021).

Implications for Conservation and Management

The changing climate and its interactions with invasive insects and native plant species pose significant challenges for conservation and management efforts. It is crucial to understand the specific ecological dynamics and vulnerabilities of native plant communities to invasive insects in the context of climate change. Implementing adaptive management strategies that consider the impacts of climate change, such as promoting habitat connectivity, supporting the establishment of native plant refuges, and integrating biological control measures, can help mitigate the negative effects of invasive insects on native plant populations (Chimweta *et. al.*, 2020; Ahissou *et. al.*, 2021).

The ecological interactions between invasive insects and native plant species are being influenced by climate change, resulting in significant consequences for biodiversity conservation and ecosystem functioning. Understanding these interactions is vital for developing effective strategies to mitigate the impacts and enhance the resilience of native plant communities. By considering the complex ecological dynamics and implementing adaptive management approaches, we can work towards maintaining the integrity and functionality of ecosystems in the face of a changing climate and invasive species threats.

2. Conclusion

The ecological interactions between invasive insects and native plant species are being significantly influenced by the changing climate, presenting substantial challenges for biodiversity conservation and ecosystem stability. Invasive insects, often introduced unintentionally or intentionally to new environments, can have detrimental impacts on native plant populations. The changing climate, with its altered phenology, range expansions, and disruptions in plant-insect interactions, exacerbates the effects of invasive insects on native plants. These impacts include increased herbivory, competitive exclusion, disruption of pollination dynamics, introduction of pathogens, and alteration of ecosystem functions. Climate change-induced phenological mismatches and range shifts further amplify the ecological disturbances caused by invasive insects, leading to declines in native plant populations, changes in community dynamics, and potential loss of ecosystem resilience. Effective management strategies, such as adaptive management approaches and integration of

biological control measures, are crucial for mitigating the impacts of invasive insects on native plant species in the context of a changing climate.

Understanding these complex ecological interactions and implementing conservation efforts can help maintain the integrity and functionality of ecosystems in the face of invasive species threats and climate change. Invasive insects, introduced to new environments either intentionally or unintentionally, can have profound impacts on native plant populations. The interactions between these two groups are influenced by a changing climate, which can lead to phenological mismatches, range expansions, altered plant-insect interactions, and feedback loops. Invasive insects can cause herbivory, competitive exclusion, disruption of pollination dynamics, introduction of pathogens, and alterations in ecosystem functions. Climate change further exacerbates these impacts, disrupting native plant communities and compromising their resilience. Effective management strategies that consider the specific ecological dynamics and vulnerabilities of native plant species are necessary to mitigate the negative effects of invasive insects. Conservation efforts should focus on understanding and preserving the integrity of ecosystems, ensuring the sustainability of native plant populations, and adapting to the challenges posed by invasive insects in the context of a changing climate.

References:

- Ahissou, B. R., Sawadogo, W. M., Bokonon-Ganta, A., Somda, I., & Verheggen, F. (2021). Integrated pest management options for the fall armyworm *Spodoptera frugiperda* in West Africa: Challenges and opportunities. A review. *Biotechnologie, Agronomie, Société et Environnement*, 25.
- Alfonzetti, M., Doleac, S., Mills, C. H., Gallagher, R. V., & Tetu, S. (2023). Characterizing Effects of Microbial Biostimulants and Whole-Soil Inoculums for Native Plant Revegetation. *Microorganisms*, 11(1), 55.
- Alfred Daniel, J., Ashok, K., Pavithran, S., & Ranjith, M. (2020). A review on invasive insect pests in India and their predators and parasitoids. *Journal of Experimental Zoology India*, 23(2), 987-1006.
- Avanesyan, A., Sutton, H., & Lamp, W. O. (2021). Choosing an effective PCR-based approach for diet analysis of insect herbivores: a systematic review. *Journal of Economic Entomology*, 114(3), 1035-1046.
- Bauer, J. T. (2012). Invasive species: "back-seat drivers" of ecosystem change? *Biological invasions*, 14, 1295-1304.
- Bhavani, B., Chandra Sekhar, V., Kishore Varma, P., Bharatha Lakshmi, M., Jamuna, P., & Swapna, B. (2019). Morphological and molecular identification of an invasive insect pest, fall army worm, *Spodoptera frugiperda* occurring on sugarcane in Andhra Pradesh, India. *Journal of Entomology and Zoology Studies*, 7(4), 12-18.
- Bisht, K., & Giri, G. S. (2019). Invasive insect pest scenario in India: a threat to biodiversity. *Journal of Entomological Research*, 43(2), 229-234.
- Cane, J. H. (2003). Exotic non-social bees (Hymenoptera: Apiformes) in North America: ecological implications. In *For non-native crops, whence pollinators of the future?* (p. 113). Entomological Society of America.
- Chakrabarti, P., Rana, S., Bandopadhyay, S., Naik, D. G., Sarkar, S., & Basu, P. (2015). Field populations of native Indian honey bees from pesticide intensive agricultural landscape show signs of impaired olfaction. *Scientific reports*, 5(1), 1-13.
- Chantawannakul, P., Ramsey, S., Khongphinitbunjong, K., & Phokasem, P. (2018). Tropilaelaps mite: an emerging threat to European honey bee. *Current opinion in insect science*, 26, 69-75.
- Chavre, B., & Patil, R. (2023). Invasive plant species of Maharashtra state: A review. *Sustainability, Agri, Food and Environmental Research*, 11.
- Chimweta, M., Nyakudya, I. W., Jimu, L., & Bray Mashingaidze, A. (2020). Fall armyworm [*Spodoptera frugiperda* (JE Smith)] damage in maize: management options for flood-recession cropping smallholder farmers. *International journal of pest management*, 66(2), 142-154.
- Crowder, D. W., & Snyder, W. E. (2010). Eating their way to the top? Mechanisms underlying the success of invasive insect generalist predators. *Biological Invasions*, 12, 2857-2876.
- Crowl, T. A., Crist, T. O., Parmenter, R. R., Belovsky, G., & Lugo, A. E. (2008). The spread of invasive species and infectious disease as drivers of ecosystem change. *Frontiers in Ecology and the Environment*, 6(5), 238-246.
- Cushman, J. H., Lortie, C. J., & Christian, C. E. (2011). Native herbivores and plant facilitation mediate the performance and distribution of an invasive exotic grass. *Journal of Ecology*, 99(2), 524-531.
- Didham, R. K., Tylianakis, J. M., Hutchison, M. A., Ewers, R. M., & Gemmell, N. J. (2005). Are invasive species the drivers of ecological change? *Trends in ecology & evolution*, 20(9), 470-474.
- Divya, J., Kalleshwaraswamy, C., Mallikarjuna, H., & Deshmukh, S. (2021). Does recently invaded fall armyworm, *Spodoptera frugiperda* displace native lepidopteran pests of maize in India. *Current Science*, 120(1358), 1358-1367.
- Drees, B. (1998). Managing red imported fire ants in wildlife areas. *Texas Fire Ant Project. On the Texas*.
- Duan, J. J., Aparicio, E., Tatman, D., Smith, M. T., & Luster, D. G. (2016). Potential new associations of North American parasitoids with the invasive Asian longhorned beetle (Coleoptera: Cerambycidae) for biological control. *Journal of economic entomology*, 109(2), 699-704.

- Elias, N. (2023). Deep Learning Methodology for Early Detection and Outbreak Prediction of Invasive Species Growth. In *Proceedings of the IEEE/CVF Winter Conference on Applications of Computer Vision* (pp. 6335-6343).
- Fortuna, T. M., Le Gall, P., Mezdoor, S., & Calatayud, P. A. (2022). Impact of invasive insects on native insect communities. *Current Opinion in Insect Science*, 51, 100904.
- Goulson, D. (2003). Effects of introduced bees on native ecosystems. *Annual Review of Ecology, Evolution, and Systematics*, 34(1), 1-26.
- Gupta, N., Verma, S. C., Sharma, P. L., Thakur, M., Sharma, P., & Devi, D. (2019). Status of invasive insect pests of India and their natural enemies. *Journal of Entomology and Zoology Studies*, 7(1), 482-489.
- Gupta, N., Verma, S. C., Sharma, P. L., Thakur, M., Sharma, P., & Devi, D. (2019). Status of invasive insect pests of India and their natural enemies. *Journal of Entomology and Zoology Studies*, 7(1), 482-489.
- Gurevitch, J., & Padilla, D. K. (2004). Are invasive species a major cause of extinctions? *Trends in ecology & evolution*, 19(9), 470-474.
- Hailu, G., Niassy, S., Bässler, T., Ochatum, N., Studer, C., Salifu, D. & Subramanian, S. (2021). Could fall armyworm, *Spodoptera frugiperda* (JE Smith) invasion in Africa contribute to the displacement of cereal stemborers in maize and sorghum cropping systems. *International Journal of Tropical Insect Science*, 41, 1753-1762.
- Hickman, J. E., Ashton, I. W., Howe, K. M., & Lerdau, M. T. (2013). The native–invasive balance: implications for nutrient cycling in ecosystems. *Oecologia*, 173, 319-328.
- Hu, J., Angeli, S., Schuetz, S., Luo, Y., & Hajek, A. E. (2009). Ecology and management of exotic and endemic Asian longhorned beetle *Anoplophora glabripennis*. *Agricultural and Forest Entomology*, 11(4), 359-375.
- Huryn, V. M. B. (1997). Ecological impacts of introduced honey bees. *The quarterly review of biology*, 72(3), 275-297.
- Jackson, M. C. (2015). Interactions among multiple invasive animals. *Ecology*, 96(8), 2035-2041.
- Jackson, M. C. (2015). Interactions among multiple invasive animals. *Ecology*, 96(8), 2035-2041.
- Jemal, A., & Hugh-Jones, M. (1993). A review of the red imported fire ant (*Solenopsis invicta* Buren) and its impacts on plant, animal, and human health. *Preventive Veterinary Medicine*, 17(1-2), 19-32.
- Johnson, P. T., Olden, J. D., Solomon, C. T., & Vander Zanden, M. J. (2009). Interactions among invaders: community and ecosystem effects of multiple invasive species in an experimental aquatic system. *Oecologia*, 159, 161-170.
- Johnson, P. T., Olden, J. D., Solomon, C. T., & Vander Zanden, M. J. (2009). Interactions among invaders: community and ecosystem effects of multiple invasive species in an experimental aquatic system. *Oecologia*, 159, 161-170.
- Kalleshwaraswamy, C. M., Asokan, R., Swamy, H. M., Maruthi, M. S., Pavithra, H. B., Hegbe, K. & Goergen, G. E. (2018). First report of the fall armyworm, *Spodoptera frugiperda* (JE Smith)(Lepidoptera: Noctuidae), an alien invasive pest on maize in India.
- Klotz, J. H., Jetter, K., Greenberg, L., Hamilton, J., Kabashima, J., & Williams, D. F. (2003). An insect pest of agricultural, urban, and wildlife areas: the red imported fire ant. In *Exotic Pests and Diseases: Biology and Economics for Biosecurity* (pp. 151-166). Iowa City: Iowa State Press.
- Kojima, Y., Toki, T., Morimoto, T., Yoshiyama, M., Kimura, K., & Kadowaki, T. (2011). Infestation of Japanese native honey bees by tracheal mite and virus from non-native European honey bees in Japan. *Microbial ecology*, 62, 895-906.
- Kumar, R., & Kumar, V. (2014). Impact of pollination by European honey bee, *Apis mellifera* L. on the yield and quality of litchi (*Litchi chinensis* Sonn.) fruits in India. *Pest Management in Horticultural Ecosystems*, 20(2), 127-132.
- LeBrun, E. G., Plowes, R. M., & Gilbert, L. E. (2012). Imported fire ants near the edge of their range: disturbance and moisture determine prevalence and impact of an invasive social insect. *Journal of Animal Ecology*, 81(4), 884-895.
- Lonighi, A. (2023). Effects of non-native species and ecological restoration on network structure and ecosystem function.
- Mallet, J. (2018). Invasive insect hybridizes with local pests. *Proceedings of the National Academy of Sciences*, 115(19), 4819-4821.
- Mani, M. (2017). Invasive insect pests and their management on tapioca (*Manihot esculenta* Crantz) in India. *Journal of Root Crops*, 43(1), 58-65.
- Marchioro, M., & Faccoli, M. (2021). Successful eradication of the Asian longhorn beetle, *Anoplophora glabripennis*, from north-eastern Italy: Protocol, techniques and results. *Insects*, 12(10), 877.
- Mills, M. D., Rader, R. B., & Belk, M. C. (2004). Complex interactions between native and invasive fish: the simultaneous effects of multiple negative interactions. *Oecologia*, 141, 713-721.
- Mills, M. D., Rader, R. B., & Belk, M. C. (2004). Complex interactions between native and invasive fish: the simultaneous effects of multiple negative interactions. *Oecologia*, 141, 713-721.
- Musolin, D. L., Kirichenko, N. I., Karpun, N. N., Aksenenko, E. V., Golub, V. B., Kerchev, I. A. & Selikhovkin, A. V. (2022). Invasive insect pests of forests and urban trees in Russia: Origin, pathways, damage, and management. *Forests*, 13(4), 521.
- Mutyambai, D. M., Niassy, S., Calatayud, P. A., & Subramanian, S. (2022). Agronomic factors influencing fall armyworm (*Spodoptera frugiperda*) infestation and damage and its co-occurrence with stemborers in maize cropping systems in Kenya. *Insects*, 13(3), 266.

- O'Neil, C. M., Guo, Y., Pierre, S., Boughton, E. H., & Qiu, J. (2023). Invasive snails alter multiple ecosystem functions in subtropical wetlands. *Science of The Total Environment*, 864, 160939.
- Paini, D. R. (2004). Impact of the introduced honey bee (*Apis mellifera*)(Hymenoptera: Apidae) on native bees: a review. *Austral ecology*, 29(4), 399-407.
- Peña, R., Schleuning, M., Miñarro, M., & García, D. (2023). Variable relationships between trait diversity and avian ecological functions in agroecosystems. *Functional Ecology*, 37(1), 87-98.
- Rathee, M., & Dalal, P. (2018). Emerging insect pests in Indian agriculture. *Indian Journal of Entomology*, 80(2), 267-281.
- Rodda, G. H., Fritts, T. H., McCoid, M. J., & Campbell III, E. W. (1999). An overview of the biology of the Brown Treesnake*(*Boiga irregularis*), a costly introduced pest on Pacific Islands.
- Rodriguez, L. F. (2006). Can invasive species facilitate native species? Evidence of how, when, and why these impacts occur. *Biological Invasions*, 8, 927-939.
- Rodriguez, L. F. (2006). Can invasive species facilitate native species? Evidence of how, when, and why these impacts occur. *Biological Invasions*, 8, 927-939.
- Sarkar, S., & Chakroborty, S. (2023). The Classification of Native and Invasive Species in North America: A Transfer Learning and Random Forest Pipeline. In *Recent Trends in Image Processing and Pattern Recognition: 5th International Conference, RTIP2R 2022, Kingsville, TX, USA, December 1-2, 2022, Revised Selected Papers* (pp. 297-307). Cham: Springer Nature Switzerland.
- Singh, S., Sharma, J. H., Udikeri, A., & Ansari, H. (2020). Invasive insects in India. In *Invasive Species-Introduction Pathways, Economic Impact, and Possible Management Options. IntechOpen*, 38, 63-74.
- Skendžić, S., Zovko, M., Pajač Živković, I., Lešić, V., & Lemić, D. (2021). Effect of climate change on introduced and native agricultural invasive insect pests in Europe. *Insects*, 12(11), 985.
- Sokame, B. M., Rebaudo, F., Malusi, P., Subramanian, S., Kilalo, D. C., Juma, G., & Calatayud, P. A. (2020). Influence of temperature on the interaction for resource utilization between Fall Armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae), and a community of lepidopteran maize stemborers larvae. *Insects*, 11(2), 73.
- Tonnang, H. E., Sokame, B. M., Abdel-Rahman, E. M., & Dubois, T. (2022). Measuring and modelling crop yield losses due to invasive insect pests under climate change. *Current Opinion in Insect Science*, 50, 100873.
- Tonnang, H. E., Sokame, B. M., Abdel-Rahman, E. M., & Dubois, T. (2022). Measuring and modelling crop yield losses due to invasive insect pests under climate change. *Current Opinion in Insect Science*, 50, 100873.
- Venette, R. C., & Hutchison, W. D. (2021). Invasive insect species: global challenges, strategies & opportunities. *Frontiers in Insect Science*, 1, 650520.
- Vinson, S. B. (1997). Insect life: invasion of the red imported fire ant (Hymenoptera: Formicidae). *American Entomologist*, 43(1), 23-39.
- Viraktamath, C. A. (2002). Alien invasive insect and mite pests and weeds in India and their management. *MICRONESICA-AGANA-*, 35, 67-83.
- Wiles, G. J., Bart, J., Beck Jr, R. E., & Aguon, C. F. (2003). Impacts of the brown tree snake: patterns of decline and species persistence in Guam's avifauna. *Conservation Biology*, 17(5), 1350-1360.
- With, K. A. (2002). The landscape ecology of invasive spread. *Conservation Biology*, 16(5), 1192-1203.
- With, K. A. (2002). The landscape ecology of invasive spread. *Conservation Biology*, 16(5), 1192-1203.
- Wojcik, D. P., Allen, C. R., Brenner, R. J., Forys, E. A., Jouvenaz, D. P., & Lutz, R. S. (2001). Red imported fire ants: impact on biodiversity.
- Wright, M. G. (2014). Biological control of invasive insect pests. In *Integrated pest management* (pp. 267-281). Academic Press.
- Zavaleta, E. S., Hobbs, R. J., & Mooney, H. A. (2001). Viewing invasive species removal in a whole-ecosystem context. *Trends in Ecology & Evolution*, 16(8), 454-459.