



Echocardiography And Multimodal Imaging Advancements In Cardiac Imaging

Hameed Ullah Baloch^{1*}, Saira Khurram², Dr. Fatima Tanveer³, Maryam Qudsia⁴, Aamir Masood⁵,
Zaibullah Khan⁶

^{1*}Department of Zoology, Wildlife and Fisheries, University of Agriculture Faisalabad, Pakistan,
Email: hameedbaloch062@gmail.com

²Subject Supervisor, Roots IVY International, Pakistan,
Email: sarakhurram28@gmail.com

³Services Institute of Medical Sciences, Pakistan,
Email: fatima_tanveer@live.com

⁴Shifa College of Medical Technology, Shifa Tameer e Millat University, Islamabad, Pakistan,
Email: qudsiastic567@gmail.com

⁵Department of Zoology, Government College University Faisalabad, Punjab, Pakistan,
Email: malikaamirmasood12@gmail.com

⁶Department of Zoology, Abdul Wali Khan University Mardan, Pakistan,
Email: zaib.zoologist@gmail.com

***Corresponding Author:** Hameed Ullah Baloch

^{*}Department of Zoology, Wildlife and Fisheries, University of Agriculture Faisalabad, Pakistan,
Email: hameedbaloch062@gmail.com

Article History	Abstract
Received: Revised: Accepted:	One of the technological sectors that has advanced and evolved the fastest in recent decades is echocardiography. As a non-invasive technique at a reasonable price, it also lends itself in the future to increasingly integrated use at the patient's bedside in any clinical approach situation, from emergencies to interventional environments, operating rooms, clinical routine, outpatient clinics, diagnostics, and monitoring of therapies. The equipment's downsizing will enable a deeper and more comprehensive integration with the clinical physical examination (clinical echocardiography), not only by cardiologists but also by the wide range of clinical specialties in medicine and surgery. This poses significant organizational and training issues to effectively incorporate multimodal imaging and ultrasound diagnostic techniques into the diagnostic and treatment paths of the various subspecialties. Additional developments in portable and miniaturized technologies are also required to achieve reliability at least on par with the highest level of equipment. This collaborative approach to multimodality imaging in cardiology may benefit from artificial intelligence.
CC License CC-BY-NC-SA 4.0	Keywords: echocardiography, multimodal imaging, ultrasound diagnostics, and cardiac imaging.

INTRODUCTION:

The development of electrocardiography, which was first documented by Waller in 1887 and made possible by Einthoven's research from its initial publication in 1901 until his discoveries were acknowledged and he was given the Nobel Prize in Medicine, can be considered the beginning of the field's first technological revolution. Electrocardiography has continued to advance, opening up new avenues for cardiological diagnosis, such as introducing contemporary sensors fitted into smartwatches (Baldassarre et al., 2022).

In the first half of the 20th century, the development of cardiac imaging was able to distinguish between the initial straightforward use of radiographic and fluoroscopic studies for cardiological diagnosis, cardiac angiography, coronary angiography, and the various applications of ultrasound, nuclear medicine imaging, computed tomography, and magnetic resonance. This development can be seen as the second technological revolution in cardiology. Multimodal imaging is another revolution that has never ceased and is still going strong (Tzolos et al., 2022).

M MODE ECHOCARDIOGRAPHY

Due to its excellent temporal resolution and single scanning direction, the M-mode (motion mode) technology is still employed in addition to other ultrasound technologies. One ultrasonic beam may switch from transmitting to receiving reflected echoes on the order of a thousand times (cycles) per second. However, two-dimensional (2D) images can only be displayed at the level of a few hundred per second. Therefore, if performed with appropriate methodology and standardization, the M-mode linear measurements thus obtained continue to represent an exact and reproducible measurement method, even in repeated examinations by different operators and other equipment (Patel, Fontana, & Ruberg, 2021).

For example, in the surgical decision of valvular diseases, with respect to each patient's linear dimensions of the left ventricle. The most complicated volumetric parameters derived from 2D or 3D images with poorer temporal resolution present variability between exams and between operators because they employ calculation methods that even measurement errors or fluctuation. The operator's decision-making in each scenario is significantly more extensive and clinically constrained. From a historical perspective, Keidel first attempted to scan the heart and measure its volumetric fluctuations in 1950 by passing an ultrasonic beam through the thorax (Agha et al., 2019).

But it wasn't until the groundbreaking research of two Swedes, Drs. Edler and Hertz were the first to employ the "ultrasonic reflectoscope" so that the heart could be thoroughly examined. As a technological foundation for later advancements, M-mode echocardiography enabled the initial studies of the anatomy and pathology of the heart and circulation. They presented a scientific film at the 3rd European Congress of Cardiology in Rome that detailed the ultrasound methods used to assess mitral stenosis, left atrial tumours, aortic stenosis, and anterior pericardial effusion. Colleagues from the United States Wild, Crawford, and Reid used the echocardiographic method to look at the heart outside the chest 1957 (Sennott & Ananthasubramaniam, 2021). Early American scholars first paid close attention to pericardial effusion. The investigation of the heart, aorta, heart valves, and their operation follows. Although the echocardiographic examination was still restricted to M mode, the 1973 publication of Harvey Feigenbaum's "Echocardiography" first edition gave the field a significant boost thanks to its wide distribution, knowledge of the physics of ultrasound, the exploration technique of the various cardiac structures, and the identification and functional evaluation of various cardiac pathologies, both congenital and acquired (Muscogiuri et al., 2022).

M-mode echocardiography has become more used in Italy since the late 1960s. The Italian Society of Cardiovascular Ultrasound (SIEC) was established in Bologna in 1973 to create the social, intellectual, technological, and practical frameworks necessary for this novel approach's advancement and clinical use. On June 17, 1974, the SIEC was formally established, and Prof. Stefano Petralia served as its inaugural president. The first National Congress of Echocardiography was organized in 1974 (Ciampi et al., 2021; Citro et al., 2020).

From Two-Dimensional To Three-Dimensional Echocardiography

The advent of 2D echocardiography, also known as "two-dimensional" or "transverse echocardiography," in the 1970s allowed for the expansion of M-mode echocardiography's capabilities, even though Wild and Reid had made the first attempts to use a "two-dimensional echoscope" as early as 1952. Cardiologists now utilize "phase array" probes instead of "linear array" probes because they can display an echocardiographic anatomical section as a fan in real-time. Having an approximate 1-2 mm spatial resolution. The probes are placed in the various intercostal spaces (also known as echocardiographic windows) during the parasternal and apical transthoracic approaches (Pergola et al., 2023).

New means of entry to the thorax, such as the subcostal or supraclavicular approach, have increased, enabling more thorough and broad imaging of the heart structures. In some circumstances, other auxiliary thoracic techniques, including the right parasternal and apical or the posterior or dorsal, have evolved through time into constructive research methods. 2D echocardiography marks a turning point and a significant technological advance in the advancement of echocardiography. Through the operator's expertise and anatomical-functional comparisons, the various 2D echocardiographic sections collected using multiple methods enable the creation of a virtual 3D depiction of the heart and its structures (Meel, 2020).

The distinct "echocardiographic windows" are constrained by the existence of the ribs and the lung, which means that 2D echocardiography also has some limits. Both hinder the transmission and penetration of ultrasound, rendering some tissues, such as the heart's atria or posterior structures, blind or challenging to examine. Networks deep in the chest result in a reduction in image quality and spatial resolution. The rate of 2D pictures might be decreased and affected by other clinical factors, such as obesity, pulmonary emphysema, lung disease, varied thoracic conformations, or even major thoracic trauma, making them subpar and occasionally unable to achieve diagnostic accuracy appropriate (Costantini et al., 2022).

The development of 3D echocardiography results from technological advancement and clinical diagnostic requirements. It began with offline reconstructions of numerous different and complementary 2D images captured from successive cardiac cycles, then progressed to "real-time" until reaching so-called "live" 3D echocardiography, which enables the representation of 2D images in 3D concurrently with the acquisition. Currently, 3D images are obtained online and in real-time for a variety of conditions and pathologies under study, including congenital heart diseases, valvular heart disease, and cardiomyopathies, and also for a better cardiac approach to the patient, from ventricular volumes to systolic function of the left ventricle. Pre-, intra-, and postoperative stages of surgery (Malik et al., 2022).

Though it offers more immediate and direct 3D spatial guidance, 3D echocardiography complements other ultrasound technologies rather than replacing them. This is because it does so at the expense of reduced temporal resolution, decreased spatial resolution, and decreased tissue characterization (Smiseth et al., 2022).

Strain Imaging, Doppler Ultrasonography, And Stress Rate

The morphology and morphofunctional characteristics of the heart can be studied using 2D and 3D echocardiography, but this technology cannot reveal the dynamics of intracardiac flows. Doppler ultrasound, also known as echo-Doppler, is a complementary technique that allows for the visualization and measurement of valve regurgitation and the possibility of studying intracavitary flows and measuring hemodynamic parameters like speed, direction, and flow regime. Based on the Doppler phenomenon (also known as the Doppler shift), which was first noticed by the Austrian Christian Johann Doppler (1803–1853) (De Lio et al., 2023).

Cardiologists measure the Doppler impact on the blood flowing through the heart. As blood flows toward the transducer, the frequency of ultrasound emitted by the device and reflected by the red blood cells they carry rises, and vice versa, as blood flows away. Because these frequency variations, also known as Doppler shifts, fall within the range of audible sound, they can be perceived as traces or spatial representations of the flow. Japanese scientist Satomura's attempt to investigate the heart's motion in 1956 is credited as the first use of Doppler ultrasound to research the organ. Doppler ultrasound was explicitly created to examine intravascular flows whose speed, strength, and direction were observed in Japan (Velaga et al., 2022).

Using the modified Bernoulli equation, it is now possible to estimate intracavitary pressures and transvalvular gradients from the measurement of velocities. These measurements are closely correlated with those made inhumanely with cardiac catheterization. The development of Doppler techniques in the 1980s allowed for the visualization of blood flow, represented using colour maps in the context of 2D images or M-mode plots. While multiple colour brightness scales spatially represent the average flow velocity, red typically denotes the direction of flow approaching the transducer and blue the receding tip (Beltrami, Dei, & Milli, 2022).

The introduction of the velocity variance representation, which determines a dispersion of colours when the flow exhibits a turbulent regime, also makes it feasible to distinguish between laminar and turbulent flow. One of the significant applications of echo-Doppler technologies in its various pulsed, continuous, and colour modes is evaluating valvular disease, which facilitates and improves the patient's ability to make clinical decisions. Every valve. Applied with the proper approach and standardization where necessary. Diastolic function, ventriculo-arterial coupling, non-invasive ultrasound hemodynamic evaluation using ultrasound, and vascular ultrasound are additional specific fields for echo-Doppler study in diverse disorders (Dorobantu et al., 2019). Since the late 1980s, researchers' focus has also broadened to include the study of the low-speed movements of the heart's valves and walls using Doppler ultrasound, which necessitates the use of specific and unique technologies. Of blood flows, using "low pass" filters that allow larger blood flow velocities to be blocked out.

This novel method, also known as tissue Doppler (DTI), "tissue speed imaging," or "Doppler myocardial imaging," allows for the measurement of the relative speeds of the various myocardial segments under pathological and ordinary circumstances, including those with global or segmental involvement, such as ischemic heart disease (Gupta-Malhotra, Schaaf, & Kutty, 2019).

DTI technologies, which allow measuring the "strain" or systolic deformation of the myocardial segments and the "strain rate" or deformation rate of the same features, can also be used to quantify regional or zonal systolic function. However, like the Doppler approach, the DTI method for analyzing deformation and strain rate necessitates an ideal alignment of the ultrasound beam with the direction of the phenomena being examined. If not, the measurements might not be accurate. By eschewing the Doppler method and instead using software that directly analyzes the representation of the "motes" (reflective components of the myocardium) or set of pixels that create the 2D images of the myocardium, new technological advancements now make it possible to measure stress and strain rate (Lin et al., 2022).

Follow the symbolic patterns of myocardial walls in grayscale or even 3D echocardiographic pictures between diastole and systole (speckle tracking). Using this technology, you can determine the myocardium's global or zonal, longitudinal, radial, and circumferential deformation values and its deformation rate. This approach appears to have great promise for population investigations, as it has the potential to detect subtle and early changes in both global and local systolic cardiac function. Its value in the daily decision-making procedures applied to each unique case in the broadest clinical circumstances is still debatable (Gudenkauf et al., 2022).

Doppler Echocardiography, Two-Dimensional, And Three-Dimensional Echocardiography In Transesophageal Approach

Transesophageal echocardiography, made possible by the probe's position in the esophagus, has created a new window for posterior heart vision, overcoming the drawbacks and restrictions of the transthoracic technique in both 2D and 3D echo. Since the transducer is closer to the heart from this vantage point, it can perform ultrasound operations at higher frequencies and with superior resolution. When 2D and then 3D echo and the use of transducers that can be positioned in various planes (biplanar and multiplanar) were introduced, transesophageal echocardiography launched in the M-mode version in the 1970s became widely used (Baggish et al., 2020).

Due to the minor inconveniences associated with its semi-invasive nature, this method is only used in specific circumstances with predetermined indications, mainly when other transthoracic windows provide insufficient or conflicting information. This new technology is always viewed as a supplement to other ultrasonic techniques rather than a replacement (Meucci, Ancona, Sanz, & Zamorano, 2022).

Intracavitary And Intracoronary Echocardiography, Contrast Ultrasound, And Interventional Echocardiography

When Gramiak and Shah used the M-mode echocardiographic technique to visualize the aortic root by injecting indocyanine green into the left ventricle during cardiac catheterization, contrast echocardiography (or contrast echocardiography) was first developed. Initially, indocyanine green, saline solution, glucose solution, or the patient's blood were employed as contrast media. The suitable heart structures were obscured by injecting these contrast materials into a peripheral vein. When right-to-left shunts were present or when the opacifying chemical was injected directly into the left heart's bloodstream during cardiac catheterization, the left heart might be reached (Figure 1) (Young et al., 2023).

These earliest medications relied on the microbubbles created when the drug was quickly given intravenously to achieve its contrast echo effect. However, the created microbubbles were frequently thick, irregular in size, unstable, and transient. Because of this, if they were injected into a peripheral vein, the pulmonary filter prevented them from entering the left circulation. Numerous studies and research projects enabled the gradual development of contrast agents with smaller, more uniform, and more stable microbubbles. These characteristics are required to overcome the pulmonary circulation and opacify the left ventricular cavity, ultimately facilitating perfusion (Chareonthaitawee & Gutberlet, 2023).

This was made possible partly by the technological introduction of the second harmonic. Myocardial. It can also be seen. Our understanding of intracardiac, coronary, and vascular circulation, as well as the perfusion and microcirculation of myocardial tissue, has substantially increased thanks to this technique. It is now only used for specific indications and investigations since it is a sophisticated, "time-consuming" method with the potential for artifacts and technical and technological restrictions that limit its widespread usage in the clinic. Therapeutic and diagnostic disciplines (Marwick et al., 2022).

The ability to position ultrasound transducers at the tip of catheters that can be inserted percutaneously (by puncturing a vein or artery) has also been made possible by the miniaturization of ultrasound transducers. These

catheters are then guided along the vascular tree to the interior of the cardiac cavities and vessels, including the coronary arteries (intracardiac or intravascular ultrasound) (Rudzinski et al., 2022).




Figure 1 shows a 68-year-old patient with dissection of the ascending aorta's M-mode echocardiography at the aortic level that was taken in the catheterization room during an aortography. The beginning of the manual injection of contrast medium (Renographin 76) is denoted in A and B by vertical arrows. The anterior false lumen (A.L.) in A and the posterior false lumen (P.L.) in B are separated by microbubbles. Aorta: (A); left atrium: (L.A.).

This kind of technological development has also made it possible to apply infrared light sources at the tip of the catheters, enabling the creation of optical coherence tomography (OCT) images with a limited depth penetration capability but providing extremely detailed sectional images of the vessel and coronary walls, comparable to echocardiographic images. Another area of application for ultrasound in cardiology that is expanding continuously and exponentially is the guidance and assessment of procedures in interventional hemodynamics and cardiac surgery (Figures 2 and 3) (Alanís-Naranjo et al., 2023).

In Various Clinical Diagnostic Circumstances And Pathological Conditions, Echocardiography

The methodology of echocardiography includes the stratification of atherosclerotic risk, athletes, complications of myocardial infarction, stress echocardiography, valvular diseases, myocardial diseases, and the pericardium, among other physiological and pathological conditions (Evangelista et al., 2023).

Anatomy-Functional Objective-Specific Echocardiography And Echocardiography Using Standard Or Miniaturized And Handheld Equipment

Increasingly portable ultrasound machines have been created due to technological advancements and the gradual miniaturization of echocardiographic equipment. These machines can be used at the patient's bedside by anyone, not just cardiologists, to complete an objective clinical examination through a study quickly. Anatomical-functional focused, confined to particular cardiac chambers or components, and allowing the gathering of data that is frequently crucial for "clinical decision-making." Its extensive use is constrained by organizational and technological issues (Bois, Anand, & Anavekar, 2019).

This is due to a variety of factors, including small screens with insufficient resolution, advanced ultrasonic techniques that are not always available on the specific device, challenges with recording and documenting images, integration challenges into clinical reports, a need for frequent in-depth analysis followed by a complete standard echocardiogram, a need for specialized training and organization of diagnostic and therapeutic procedures and protocols, and the possibility of a higher nadir (Mada, Mada, Stef, Molnar, & Rosianu, 2022).

Cardiac Multimode Imaging

The proliferation of imaging technologies has made it possible to delve into many anatomical-functional and cardiovascular pathological features, frequently providing information complementary to echocardiography in various cardiac diseases. Although these technologies, such as nuclear cardiology, cardiac magnetic resonance,

and computed tomography, are available, they are less widely dispersed throughout the territory. They are more difficult to access, even when they are, making them less useful, particularly in emergencies (Donal et al., 2019).

All of this has, up to now, encouraged the use of it that is not necessarily reasonable and appropriate, with a propensity toward overuse in institutions with particular programs, protocols, and organizational tools but also toward underuse in circumstances where it is unavailable or challenging to get, simply due to scheduling, seasonal, or vacation constraints. The challenges brought on by the requirement for multi-specialized training of new intermediate or hybrid professional figures between cardiology and radiology, outside of cardiology, are added to this. The co-responsible involvement of both professional figures in the creation and implementation of diagnostic protocols and in the interpretation and reporting of tests may provide a solution to this challenge (Anghel et al., 2020).

Implementing this excellent possibility for a shared interdisciplinary approach to cardiovascular disorders in many institutions can be challenging. In this setting, artificial intelligence can help advance and solidify a cardiology imaging strategy that is becoming increasingly integrated (Galea et al., 2020).




Figure 2 shows two-dimensional (A, E, I) and three-dimensional (other panels) transesophageal echocardiographic monitoring of a transcatheter mitral clip application process. The arrow in B points to the location where the catheter tip used to puncture the interatrial septum pushes into the oval fossa, taking on the shape of a tent. The arrow in C denotes the separation between the mitral valve (MV) and the tenting (T). D illustrates the guidewire passing through the atrial septum (AIS). In E and F, the guiding catheter is displayed. The clip delivery system (red arrow) is directed at the mitral valve in panels G–N. The clip appears closed in M, and in N, it is open (Maloku et al., 2023).




Figure 3 shows the placement of a percutaneous mitral prosthetic ring (Valtech Cardioband System) followed by a fluoroscopic inspection (top) and a three-dimensional transesophageal echo (middle and bottom left). To Available online at: <https://jazindia.com>

achieve the native mitral annulus' ideal dimensions for a decrease in mitral regurgitation, as shown by the colour Doppler (bottom right), the stepwise implantation of the prosthetic ring anchors is underlined (Ranganath et al., 2020).

CONCLUSIONS:

One of the technological sectors that has advanced and evolved the fastest in recent decades is echocardiography. It is also appropriate for the future because it is a non-intrusive and reasonably priced method from emergencies, intervention settings, operating rooms, clinical routine, outpatient consultations, diagnosis, prognosis, and therapy monitoring to an increasingly integrated clinical use at the patient's bedside. It also lends itself to a deeper and more complementary integration with the objective clinical examination (clinical echocardiography), not just by the cardiologist but also by the variety of medical and surgical clinical specialties, thanks to the equipment's ongoing miniaturization.

This poses significant organizational and training problems to effectively incorporate multimodal imaging and ultrasound diagnostic tools into the various specialized diagnostic and therapy paths. Miniaturized and portable technologies also need to keep developing to reach a degree of dependability equal to that of mobile devices. Several premium standard products. The multidisciplinary approach to multimodal imaging in cardiology will benefit from using artificial intelligence in this scenario.

REFERENCES:

1. Agha, A. M., Lopez-Mattei, J., Donisan, T., Balanescu, D., Iliescu, C. A., Banchs, J., . . . Gladish, G. (2019). Multimodality imaging in carcinoid heart disease. *Open heart*, 6(1), e001060.
2. Alanís-Naranjo, J. M., Galeana-Abarca, C., Rentería-Galicia, M. A., Arroyo-Martínez, E. A., Loredó-Alanís, M. A., & Rodríguez-López, T. (2023). Left atrial myxoma: the importance of cardiovascular imaging in the long-term follow-up of breast cancer survivors.
3. Anghel, L., Stătescu, C., Șerban, I.-L., Măranducă, M. A., Butcovan, D., Clement, A., . . . Sascău, R. (2020). The Advantages of New Multimodality Imaging in Choosing the Optimal Management Strategy for Patients with Hypertrophic Cardiomyopathy. *Diagnostics*, 10(9), 719.
4. Baggish, A. L., Battle, R. W., Beaver, T. A., Border, W. L., Douglas, P. S., Kramer, C. M., . . . Singh, T. K. (2020). Recommendations on the use of multimodality cardiovascular imaging in young adult competitive athletes: a report from the American Society of Echocardiography in Collaboration with the Society of Cardiovascular Computed Tomography and the Society for Cardiovascular Magnetic Resonance. *Journal of the American Society of Echocardiography*, 33(5), 523-549.
5. Baldassarre, L. A., Ganatra, S., Lopez-Mattei, J., Yang, E. H., Zaha, V. G., Wong, T. C., . . . Deswal, A. (2022). Advances in multimodality imaging in cardio-oncology: JACC state-of-the-art review. *Journal of the American College of Cardiology*, 80(16), 1560-1578.
6. Beltrami, M., Dei, L.-L., & Milli, M. (2022). The Role of the Left Atrium: From Multimodality Imaging to Clinical Practice: A Review. *Life*, 12(8), 1191.
7. Bois, J. P., Anand, V., & Anavekar, N. S. (2019). Detection of inflammatory aortopathies using multimodality imaging. *Circulation: Cardiovascular Imaging*, 12(7), e008471.
8. Chareonthaitawee, P., & Gutberlet, M. (2023). Clinical Utilization of Multimodality Imaging for Myocarditis and Cardiac Sarcoidosis. *Circulation: Cardiovascular Imaging*, 16(1), e014091.
9. Ciampi, Q., Antonini-Canterin, F., Barbieri, A., Barchitta, A., Benedetto, F., Cresti, A., . . . Pepi, M. (2021). Remodeling of activities of Italian echocardiographic laboratories during the coronavirus disease 2019 lockdown: the SIECoVID study. *Journal of Cardiovascular Medicine*, 22(7), 600-602.
10. Citro, R., Okura, H., Ghadri, J. R., Izumi, C., Meimoun, P., Izumo, M., . . . Kagiya, N. (2020). Multimodality imaging in takotsubo syndrome: a joint consensus document of the European Association of Cardiovascular Imaging (EACVI) and the Japanese Society of Echocardiography (JSE). *European Heart Journal-Cardiovascular Imaging*, 21(11), 1184-1207.
11. Costantini, P., Perone, F., Siani, A., Groenhoff, L., Muscogiuri, G., Sironi, S., . . . Guglielmo, M. (2022). Multimodality Imaging of the Neglected Valve: Role of Echocardiography, Cardiac Magnetic Resonance and Cardiac Computed Tomography in Pulmonary Stenosis and Regurgitation. *Journal of Imaging*, 8(10), 278.
12. De Lio, F., Andreis, A., De Lio, G., Bellettini, M., Pidello, S., Raineri, C., . . . Imazio, M. (2023). Cardiac imaging for the prediction of sudden cardiac arrest in patients with heart failure. *Heliyon*.

13. Donal, E., Delgado, V., Bucciarelli-Ducci, C., Galli, E., Haugaa, K. H., Charron, P., . . . Galderisi, M. (2019). Multimodality imaging in the diagnosis, risk stratification, and management of patients with dilated cardiomyopathies: an expert consensus document from the European Association of Cardiovascular Imaging. *European Heart Journal-Cardiovascular Imaging*, 20(10), 1075-1093.
14. Dorobantu, L., Ticulescu, R., Greavu, M., Dermengiu, A., Alexandrescu, M., & Trofin, M. (2019). Current management and surgical advances in patients with hypertrophic obstructive cardiomyopathy. *Kardiologia Polska (Polish Heart Journal)*, 77(9), 829-836.
15. Evangelista, A., Sitges, M., Jondeau, G., Nijveldt, R., Pepi, M., Cuellar, H., . . . Dweck, M. R. (2023). Multimodality imaging in thoracic aortic diseases: a clinical consensus statement from the European Association of Cardiovascular Imaging and the European Society of Cardiology working group on aorta and peripheral vascular diseases. *European Heart Journal-Cardiovascular Imaging*, 24(5), e65-e85.
16. Galea, N., Bandera, F., Lauri, C., Autore, C., Laghi, A., & Erba, P. A. (2020). Multimodality imaging in the diagnostic work-up of endocarditis and cardiac implantable electronic device (CIED) infection. *Journal of Clinical Medicine*, 9(7), 2237.
17. Gudenkauf, B., Hays, A. G., Tamis-Holland, J., Trost, J., Ambinder, D. I., Wu, K. C., . . . Sharma, G. (2022). Role of multimodality imaging in the assessment of myocardial infarction with nonobstructive coronary arteries: beyond conventional coronary angiography. *Journal of the American Heart Association*, 11(1), e022787.
18. Gupta-Malhotra, M., Schaaf, W., & Kutty, S. (2019). A primer on multimodal imaging and cardiology-radiology congenital heart interface. *Children*, 6(4), 61.
19. Lin, Y., Wu, W., Gao, L., Ji, M., Xie, M., & Li, Y. (2022). Multimodality imaging of benign primary cardiac tumor. *Diagnostics*, 12(10), 2543.
20. Mada, C., Mada, R. O., Stef, A., Molnar, A., & Rosianu, H. S. (2022). An unusual complication of myocardial infarction. *Medical Ultrasonography*, 24(4), 496-498.
21. Malik, N., Mukherjee, M., Wu, K. C., Zimmerman, S. L., Zhan, J., Calkins, H., . . . Tandri, H. (2022). Multimodality imaging in arrhythmogenic right ventricular cardiomyopathy. *Circulation: Cardiovascular Imaging*, 15(2), e013725.
22. Maloku, A., Hamadanchi, A., Bargenda, S., Richter, M., Möbius-Winkler, S., Schulze, C., & Franz, M. (2023). Management of giant left atrial thrombus late after Transcatheter Mitral Valve-in-ring Replacement using a Transcatheter Aortic Valve.
23. Marwick, T. H., Gimelli, A., Plein, S., Bax, J. J., Charron, P., Delgado, V., . . . Maurovich-Horvat, P. (2022). Multimodality imaging approach to left ventricular dysfunction in diabetes: an expert consensus document from the European Association of Cardiovascular Imaging. *European Heart Journal-Cardiovascular Imaging*, 23(2), e62-e84.
24. Meel, R. (2020). Incorporating multimodality imaging in training in the South African context. *SA Heart*, 17(3), 381-383.
25. Meucci, F., Ancona, F., Sanz, A. P., & Zamorano, J. (2022). Echocardiographic guidance in transcatheter structural cardiac interventions. *EuroIntervention*, 17(15), 1205-1226.
26. Muscogiuri, G., Volpato, V., Cau, R., Chiesa, M., Saba, L., Guglielmo, M., . . . Dell'Aversana, S. (2022). Application of AI in cardiovascular multimodality imaging. *Heliyon*.
27. Patel, R. K., Fontana, M., & Ruberg, F. L. (2021). Cardiac amyloidosis: multimodal imaging of disease activity and response to treatment. *Circulation: Cardiovascular Imaging*, 14(6), e009025.
28. Pergola, V., Cameli, M., Mushtaq, S., Mattesi, G., D'Andrea, A., Guaricci, A. I., . . . Motta, R. (2023). Multimodality Imaging in Advanced Heart Failure for Diagnosis, Management and Follow-Up. A Comprehensive Review.
29. Ranganath, P., Moore, A., Guerrero, M., Collins, J., Foley, T., Williamson, E., & Rajiah, P. (2020). CT for pre-and postprocedural evaluation of transcatheter mitral valve replacement. *Radiographics*, 40(6), 1528-1553.
30. Rudzinski, P. N., Leipsic, J. A., Schoepf, U. J., Dudek, D., Schwarz, F., Andreas, M., . . . Burt, J. R. (2022). CT in transcatheter-delivered treatment of valvular heart disease. *Radiology*, 304(1), 4-17.
31. Sennott, J., & Ananthasubramaniam, K. (2021). Multimodality imaging approach to cardiac amyloidosis: Part 2. *Heart failure reviews*, 1-16.
32. Smiseth, O. A., Morris, D. A., Cardim, N., Cikes, M., Delgado, V., Donal, E., . . . Gimelli, A. (2022). Multimodality imaging in patients with heart failure and preserved ejection fraction: an expert consensus document of the European Association of Cardiovascular Imaging. *European Heart Journal-Cardiovascular Imaging*, 23(2), e34-e61.

33. Tzolos, E., Kwiecinski, J., Berman, D., Slomka, P., Newby, D. E., & Dweck, M. R. (2022). Latest advances in multimodality imaging of aortic stenosis. *Journal of Nuclear Medicine*, 63(3), 353-358.
34. Velaga, J., Liew, C., Choo Poh, A. C., Lee, P. T., Lath, N., Low, S. C., & Bharadwaj, P. (2022). Multimodality imaging in the diagnosis and assessment of cardiac amyloidosis. *World Journal of Nuclear Medicine*, 21(03), 173-183.
35. Young, K. A., Raoult, T., Leccisotti, L., Gerber, B., Chareonthaitawee, P., & Gheysens, O. (2023). Multimodality imaging for diagnosis, risk stratification and treatment monitoring of cardiac sarcoidosis.