



Crossing Numbers of the Cartesian Product of the Double Triangular Snake Graphs With Path P_m .

Pathak Manojkumar Vijaynath^{1*}, Dr. Nithya Sai Narayana²

^{1*}VPM'S Kannada high school and Junior college, Mithagar road, Mulund(E), Mumbai-81.

manojkumarpatak82v@gmail.com

²Department of Mathematics, University of Mumbai, Vidyanagari Kalina Santacruz East Mumbai-400098.

narayana nithya@yahoo.com

***Corresponding Author:** Pathak Manojkumar Vijaynath

¹VPM'S Kannada high school and Junior college, Mithagar road, Mulund(E), Mumbai-81.

manojkumarpatak82v@gmail.com

CC License CC-BY-NC-SA 4.0	Abstract The crossing number $Cr(G)$ of a graph G is the least number of edge crossings in all possible good drawings of G in the plane. Join and Cartesian products of graphs have many interesting graph-theoretical properties. In this paper, we evaluate the crossing number of the Cartesian product of double triangular snake graph DT_2 with the path P_m. In this paper, we proved $Cr(DT_2 \times P_m) = 6(m - 2)$, for $m \geq 2$ where Cr denotes the crossing number. Keywords: Crossing number, Cartesian product, Join product, Double triangular snake graph, Path, cycle. Mathematics Subject Classification (2010): 68R10; 05C10; 05C62
--------------------------------------	---

Introduction

The crossing number of a graph G is the least possible number of edge crossings in any drawing of G in the plane. It is denoted by $Cr(G)$. A drawing of a graph is good if every two edges have at most one point in common, which is either a common end vertex or a crossing. We denote the number of crossings of a graph G in a drawing τ in the plane by $Cr_\tau(G)$. For two subgraphs H and H' of G such that $E(H) \cap E(H') \neq \emptyset$, $Cr_\tau(H, H')$ denotes the edge crossings between $E(H)$ and $E(H')$ in the drawing τ . The union of the two graphs F and K , denoted by $F \cup K$ and it is the graph with vertex set $V(F) \cup V(K)$ and edge set $E(F) \cup E(K)$. The join of the two graphs F and K , denoted by $F + K$, is the graph obtained from $F \cup K$ by joining every vertex of F to every vertex of K .

The Cartesian product of the two graphs F and K , denoted by $F \times K$, is the graph whose vertex set is $V(F) \times V(K)$ and the edges are of the form $(a, b)(m, n)$ where either $a = m$ and $bn \in E(K)$ or $b = n$ and $am \in E(F)$.

Let P_m denote the path on m vertices of length $(m-1)$. There are few results with $Cr(G \times P_m)$, where G is a graph on 7 vertices. Finding the value of $Cr(G \times P_m)$ has been investigated in [1], [2], [3], [5], [6], [7], [8], [9], [10]. In this paper, we evaluated the exact value of $Cr(DT_2 \times P_m)$ for $m \geq 2$.

1 Crossing numbers of the Cartesian prod- uct of Double Triangular snake graph DT_2 with P_m .

Definition 2.1. (Double Triangular Snake graph DT_p) A graph DT_p with p blocks is obtained from a path on vertices $\alpha_0, \alpha_1, \alpha_2, \dots, \alpha_p$ and by joining α_i and α_{i+1} to two new vertices α_{p+1+i} and α_{2p+1+i} for $i=0,1,2,\dots,p-1$




Figure 1: (Double Triangular snake with 5 blocks)

Drawing of $DT_2 \times P_m$:

For $m \geq 2$, Graph $DT_2 \times P_m$ has $7m$ vertices and $(17m - 7)$ edges that are the edges in the m copies of graph DT_2 and in the 7 paths of m vertices.

Notation:

- (1) $(DT_2)^i$: i^{th} copy of Double Triangular Snake graph DT_2 .
- (2) $(E_2)^i$: Edges which join vertices of $(i + 1)^{\text{th}}$ copy of Double Tri- angular Snake graph DT_2 with corresponding vertices of i^{th} copy of the graph DT_2 .
- (3)




Figure 2: ($DT_2 \times P_m$)

Remark 2.1. $Cr(DT_2 \times P_2) = 0$.




Figure 3: ($DT_2 \times P_2$)

Lemma 2.1. Let $Z(n, m) = \left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n-1}{2} \right\rfloor \left\lfloor \frac{m}{2} \right\rfloor \left\lfloor \frac{m-1}{2} \right\rfloor$

- (i) $\text{Cr}(K_{n,m}) = Z(n, m)$ where $\min\{m, n\} \leq 6$,
(ii) $\text{Cr}(K_{n,m}) \leq Z(n, m)$ for $m, n \in \mathbb{N}$.

Theorem 2.1. $\text{Cr}_\tau(DT_2 + 2K_1) = 6$.

Proof. $DT_2 + K_1$ is planar. Consider a good drawing of the graph $DT_2 + 2K_1$ given in **fig. 4**.




Figure 4: $(DT_2 + 2K_1)$

From fig. 4 we can conclude that,

$$\text{Cr}_\tau(DT_2 + 2K_1) \leq 6 \quad (1)$$

Let us prove the otherway inequality.

Any drawing τ of the graph $DT_2 + 2K_1$ contains a sub drawing of $K_{3,6}$. Thus $\text{Cr}_\tau(DT_2 + 2K_1) \geq \text{Cr}_\tau(K_{3,6}) = 6$.

$$\text{Cr}_\tau(DT_2 + 2K_1) \geq 6 \quad (2)$$

form **eq. (1)** and **eq. (2)**, we can conclude that,

$$\text{Cr}_\tau(DT_2 + 2K_1) = 6$$

Lemma 2.2. For any good drawing of the graph $DT_2 \times P_2$, let y be a vertex such that $y \notin V(DT_2)$ and graph S obtained by joining all vertices of one copy of the graph DT_2 with the vertex y then $\text{Cr}_{\tau_1}(S) = 6$.

Proof. Consider a good drawing of the graph S , as shown in **fig. 5**.




Figure 5: ((Drawing of the graph S))

Thus from **fig. 5** we can conclude that,

$$\text{Cr}_{\tau_1}(S) \leq 6.$$

Now we have to prove other way inequality. By contracting all vertices and edges of another copy of the graph DT_2 in any drawing τ_1 of graph S , we get a contraction of subdrawing of the graph $DT_2 + 2K_1$. Thus,

$$\text{Cr}_{\tau_1}(S) \geq \text{Cr}_{\tau_1}(DT_2 + 2K_1) = 6.$$

$$\text{Cr}_{\tau_1}(S) \geq 6. \text{ Hence proved.}$$

Theorem 2.2. $Cr_t(DT_2 \times P_3) = 6$.

Proof. Consider a good drawing of the graph $DT_2 \times P_3$ as shown in **fig. 6**.




Figure 6: $(DT_2 \times P_3)$

Thus from **fig. 6** we can conclude that,

$$Cr_t(DT_2 \times P_3) \leq 6$$

Now we have to prove otherway inequality. By contracting all vertices and edges of 3rd copy of the graph DT_2 in any drawing τ of the graph $DT_2 \times P_3$, we get a contraction of the graph S . Thus,

$$Cr_t(DT_2 \times P_3) \geq Cr_t(S) = 6$$

$$Cr_t(DT_2 \times P_3) \geq 6.$$

Hence proved.

Theorem 2.3. $Cr_t(DT_2 \times P_m) = 6(m - 2)$, for $m \geq 2$.

Proof. We prove this by the method of induction on m . By **theorem 2.2** result holds for $m=3$. Let us assume the result holds for less than m . Now we have to prove the result for m . In any good drawing τ of the graph $DT_2 \times P_m$, by Contracting all vertices and edges of n^{th} copy of the graph DT_2 . We get a drawing of a disjoint union of subgraphs $DT_2 \times P_{m-1}$ and S . Thus

$$Cr_t(DT_2 \times P_m) \geq Cr_t(DT_2 \times P_{m-1}) + Cr_t(S) = 6(m-3) + 6 = 6(m-2)$$

$$Cr_t(DT_2 \times P_m) \geq 6(m - 2)$$

and from the drawing of the graph $DT_2 \times P_m$ shown in **fig. 2**, we can conclude that,

$$Cr_t(DT_2 \times P_m) \leq 6(m - 2)$$

Hence proved.

2 Conclusion

In this paper, we obtained the exact crossing number of the Cartesian product of double triangular snake graph DT_2 with P_m .

3 Acknowledgment

I express my sincere gratitude to my guide Dr. Nithya Sai Narayana for her valuable guidance. I am also thankful to the Mathematics department, the University of Mumbai for selecting me as a researcher in this subject. My sincere thanks also go to my college for their support. I also thank the Editorial Board of this journal for publishing my work.

4 Funding

Not applicable

5 Data availability statement

Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

6Declarations

This declaration is not applicable.

7 Conflict of interest

There is no any conflict of interest.

References

1. Bokal, D. On the crossing numbers of Cartesian products with paths. *Journal of Combinatorial Theory, Series B* 97, 3(2007),381–384.
2. Klesc, M.” The crossing numbers of products of paths and stars with 4-vertex graphs.” *Journal of Graph Theory* 18, 6(1994), 605–614.
3. Klesc, M.” The crossing numbers of cartesian products of paths with 5-vertex graphs”. *Discrete Mathematics* 233,1-3 (2001),353–359.
4. Maria’n Kles’c’ and stefan Schro’tter,” THE CROSSING NUMBERS OF JOIN PRODUCTS OF PATHS WITH GRAPHS OF ORDERFOUR”. *Discussiones Mathematicae Graph Theory* 31(2)(2011) 321- 331.
5. Ouyang, Z., Wang, J., and Huang, Y. The crossing number of the Cartesian product of paths with complete graphs. *Discrete Mathe- matics* 328 (2014), 71–78.
6. Pathak Manojkumar Vijaynath and Dr. Nithya Sai Narayana” On the Crossing numbers of $TS_n \times P_m$ and $TS_n \times C_m$.” *Journal of Com- putational Mathematica* (2023), 2456-8686.
7. Peng, Y., and Yiew, Y. The crossing number of $p(3, 1) \times P_n$. *Dis- crete mathematics* 306, 16 (2006), 1941–1946.
8. Yiew, Y. C., Chia, G. L., and Ong, P.-H. Crossing number of the Cartesian product of prism and path. *AKCE International Journal of Graphs and Combinatorics* 0, 0 (2020), 1–7.
9. Yuan, Z., and Huang, Y. The crossing number of petersen graph $P(4, 1)$ with paths P_n . *Oper. Res. Trans* 15, 3 (2011), 95–106.
10. Zheng, W., Lin, X., and Yang, Y. The crossing number of $K_{2,m} \times P_n$, *Discrete Mathematics* 308, 24 (2008), 6639–6644.